

Davis, CA

# City Tree Resource Analysis

Analysis of Inventoried City Trees (2022)







Davis, CA

# City Tree Resource Analysis

Analysis of Inventoried City Trees (2022)



**Prepared For:**

City of Davis  
23 Russell Blvd.  
Davis, CA 95616



**Prepared By:**

Davey Resource Group, Inc.  
6005 Capistrano Avenue  
Atascadero, CA 93422  
[www.davey.com/drg](http://www.davey.com/drg)

***Acknowledgements:***

This project is funded in-part by CAL FIRE through Prop 68.

# Table of Contents

Executive Summary.....	1
Structure.....	1
Benefits.....	2
Management & Investment.....	2
Introduction.....	5
Resource Structure.....	7
Species Diversity.....	7
Importance Value.....	9
Canopy Cover.....	10
Relative Age Distribution.....	11
Tree Condition & Relative Performance.....	13
Relative Performance Index.....	13
Replacement Value.....	15
Resource Benefits.....	17
Air Quality.....	17
Deposition, Interception, & Avoided Pollutants.....	18
Atmospheric Carbon Dioxide Reductions.....	21
Stormwater Runoff Reductions.....	23
Energy Savings.....	26
Electricity & Natural Gas Reductions.....	26
Aesthetic, Property Value, & Socioeconomic Benefits.....	26
Calculating Tree Benefits.....	27
Annual Benefits of Most Prevalent Species.....	27
Net Annual Benefits.....	30
Annual Investment & Benefit Offset.....	30
Pest and Pathogen Threats.....	31
Pest Management.....	34
Conclusion.....	35
Appendix A: References.....	37
Appendix B: Methods.....	41
i-Tree <i>Eco</i> Model and Field Measurements.....	41
Definitions and Calculations.....	41
Appendix C: Tables.....	43

## Figures

Figure 1: Annual Benefits from the City Tree Resource.....	2
Figure 2: Species Diversity in Davis' City tree resource .....	8
Figure 3: Community Tree Inventory Relative Age Distribution.....	11
Figure 4: Relative Age Distribution of Davis' Top 10 Most Prevalent Species.....	12
Figure 5: Tree Condition .....	13
Figure 6: Annual Air Pollution Benefits .....	17
Figure 7: Top 5 Species for Air Pollution Removal Benefits.....	20
Figure 8: Top 5 Species for Carbon Benefits.....	22
Figure 9: How Trees Impact Stormwater.....	23
Figure 10: Top 5 Species for Stormwater Benefits.....	24
Figure 11: Summary of Annual Benefits for Most Prevalent Species .....	28
Figure 12: Annual Environmental Benefits.....	30

## Tables

Table 1: Population Summary of Most Prevalent Species.....	7
Table 2: Importance Value (IV) of Prevalent Species in Davis (Representing > 1%) .....	10
Table 3: Relative Performance Index of Most Prevalent Species .....	14
Table 4: Species That May Be Underused (based on RPI and age distribution) .....	15
Table 5: Replacement Value for Most Prevalent Species.....	16
Table 6: Adverse Health Incidents Avoided Due to Changes in Pollutant Concentration Levels and Economic Values.....	18
Table 7: Annual Air Pollution Removal Benefits .....	18
Table 8: Annual Air Quality Benefits by Most Prevalent Species .....	20
Table 9: Annual Carbon Sequestration Benefits by Most Prevalent Species.....	22
Table 10: Stormwater Benefits by Most Prevalent Tree Species.....	24
Table 11: Summary of Annual Benefits of Most Prevalent Species .....	29
Table 12: Quantifiable Benefits and Investments.....	30
Table 13: Pest & Pathogen Threats to Davis .....	32
Table 14: Botanical and Common Names of Tree Species in Davis' City tree resource.....	43
Table 15: Population Summary for All Species .....	48
Table 16: Importance Value (IV) for All Tree Species .....	53
Table 17: Condition and RPI for All Tree Species.....	58
Table 18: Annual Benefits for All Species.....	63

## Executive Summary

City trees (also called community trees or street trees) are trees in the public rights-of-way, including trees along streets, in medians, and in parks. While they are only a subset of the entire community forest, City trees play a vital role in the City of Davis and are an important contribution to the larger urban forest in the City. They provide numerous tangible and intangible benefits to residents, employees, visitors, and neighboring communities. The City recognizes that trees are a valued resource, a critical component of the urban infrastructure, and part of the community's identity.

The City of Davis contracted with Davey Resource Group, Inc. (DRG) to use the City's community tree inventory data in conjunction with i-Tree *Eco* benefit-cost modeling software to develop a detailed and quantified analysis of the current structure, function, benefits, and value of the City's community tree resource. This report details the results of that analysis. It is important to note that this analysis does not include private trees not within the City's inventory.

## Structure

A structural analysis is the first step towards understanding the benefits provided by City trees, as well as their management needs. Davis' City tree inventory includes 30,692 trees and 407 available planting sites. Considering species diversity, age distribution, condition, canopy coverage, and replacement value, DRG determined that the following information characterizes the City tree inventory:

- 207 unique tree species
- *Platanus x acerifolia* (London planetree, 7.8%) is the most common species, followed by *Pistacia chinensis* (Chinese pistache, 6.6%), and *Lagerstroemia indica* (common crapemyrtle, 6.2%)
- 36% of trees are less than 8 inches in diameter (DBH<sup>1</sup>) and 9.8% of trees are larger than 24 inches in diameter, indicating a well-established age distribution
- Based on tree inventory data (2018), 93.3% of City trees are in fair or better condition
- City trees provide an estimated 21.2% canopy cover
- City trees have stored more than 16,158 tons of carbon (CO<sub>2</sub>) in woody and foliar biomass
- To replace Davis' 30,692 City trees with trees of equivalent size, species, and condition, would cost over \$91.5 million
- 69% of Davis' City trees are susceptible to identified pests and disease threats such as polyphagous shot hole borer, defoliating moths, thousand cankers disease and Dutch elm disease

---


<sup>1</sup>DBH: Diameter at Breast Height. DBH represents the diameter of the tree when measured at 1.4 meters (4.5 feet) above ground (U.S.A. standard).

# Benefits

Many of the benefits from urban trees cannot be accurately quantified with current formulas and peer-reviewed consensus. Numerous studies indicate that urban trees have innumerable critical benefits to natural ecosystems, economies, and human health and welfare. However, i-Tree *Eco* is currently limited to quantifying the benefits from trees to air quality, stormwater runoff reduction, and carbon sequestration. In addition, this report does not portray the benefits from all of Davis' trees. Private trees are not included in the analysis, therefore this report represents only the benefits received from City trees.

Annually, City trees provide quantifiable benefits to the community totaling \$213,857. The average annual benefit per tree is \$6.97. These benefits include:

- 2.7 million gallons of avoided stormwater runoff, valued at \$24,552, an average of \$0.80 per tree
- 10.4 tons of air particulates removed, improving air quality, and reducing adverse health incidents for a value of \$117,423, an average of \$3.83 per tree
- 421.5 tons of carbon directly sequestered, valued at \$71,882, an average of \$2.34 per tree



**Figure 1: Annual Benefits from the City Tree Resource**

# Management & Investment

Annually, the City invests approximately \$1.6 million (\$51.41/tree, \$22.77/capita) to manage public trees. The quantifiable benefits from i-Tree *Eco* offset this investment by \$213,857, for a net investment of \$1.4 million. However, this offset amount is inarguably a conservative estimate of the true environmental and socioeconomic benefits from this vital resource, including, benefits to wildlife, property values, and public health and welfare. Additionally, when tree data includes the distance and direction from nearby buildings, i-Tree *Eco* can calculate estimated energy savings (gas and electric) resulting from the shade and protection of trees. The inventory does not currently include these metrics.

The City of Davis' tree inventory is a dynamic resource that requires continued investment to maintain and realize its full benefit potential. Trees are one of the few community assets that have the potential to increase in value with time and proper management. Appropriate and timely tree care can substantially increase lifespan and benefit yield. When trees live longer, they provide greater benefits. As individual trees mature, and aging trees are replaced, the overall value of the community forest and the amount of benefits provided grow as well. However, this vital living resource is vulnerable to a host of stressors and requires ecologically sound and sustainable best management practices to ensure a continued flow of benefits for future generations.

Although urban forest managers cannot foresee when a pest or pathogen may be introduced to the community forest, being aware of and able to identify potential threats allows them to approach management and prevention in a way that fits the community's culture and available resources. Using best management practices to prepare for and/or manage pests and pathogens can lessen the detrimental impacts they have on the community forest.


Overall, the City tree inventory is a resource in fair or better condition with a well-established age distribution. With proactive management, planning, and new and replacement tree planting, the benefits from this resource will continue to increase as young trees mature.

Based on this resource analysis, DRG recommends the following:

- Regularly inspect trees to identify and mitigate structural and age-related defects to manage risk and reduce the likelihood of tree and branch failure.
- Provide structural pruning for young trees and a routine pruning cycle for all trees.
- Increase species diversity in new and replacement tree plantings to reduce reliance on the most prevalent species and increase resilience.
  - Discourage the planting of species that are classified as invasive.
- Monitor species performance (e.g., health, structure, longevity, pest and disease resistance) and consider new, promising species in future tree plantings.
- Consider successional planting of important species, based on relative performance and relative age distributions.
- Replace trees that have been removed and increase the stocking level for optimal benefits.
- Plant large-stature species for greater benefits wherever space allows.
- Follow best management practices, when monitoring for and dealing with pests and diseases.
- Maintain and update the inventory database to include all public trees (including in open space), all available planting sites, track tree growth and condition, and consider adding distance and direction from buildings to calculate energy benefits.

With adequate protection and planning, the value of the City tree resource will continue to increase over time. Proactive management and a tree replacement plan are critical to ensuring that the community continues to receive a high level of benefits. Along with new tree installations and replacement plantings, funding for tree maintenance and inspection is necessary to preserve benefits, prolong tree life, and manage risk and public safety. Existing mature trees should be maintained and protected whenever possible since the greatest environmental benefits accrue from the continued growth and longevity of the existing canopy. Urban forest managers can take pride in knowing that City trees support the quality of life for residents and neighboring communities.





*The greatest benefits accrue from the continued growth and longevity of the existing canopy.*



# Introduction

Davis is a small city located in the Central Valley of California, roughly 15 miles west of the California State Capitol in Sacramento. The area has a reputation as a center of agriculture for its rich soil, multimillion dollar agricultural industry/commodities, and the nation's top rated agricultural college, the University of California at Davis. As a City, Davis is known for its walkable downtown, friendly community, and farmers markets which showcase the surrounding agricultural industry of the area. The City of Davis, perhaps most well known for being a bicycle-friendly community, has earned the title of the 'Bicycle Capital of America' as early as 1964 and was the first city in the United States to earn the Platinum Bicycle Friendly Community award from the League of American Bicyclists in 2006. Today, careful planning, not only for bicycle use but for growth as a whole, is seen throughout the city and evidenced by a connected network of greenbelts, dedicated bike and pedestrian paths, and bicycle tunnels and bridges.

Davis experiences a Mediterranean climate with an average of 21 inches of rainfall each year, most of which occurs in the winter and spring months. Most of the precipitation falls in the winter, with daytime temperatures that average in the 50s. In the summer, daytime temperatures in Davis average in the mid-90s. There are 267 days of sunshine each year and temperatures do not typically drop below freezing (Sperling's Best Places, n.d.).

Urban trees play an essential role in the community providing many benefits, tangible and intangible, to residents, visitors, and neighboring communities. Research demonstrates that healthy urban trees can improve the local environment and lessen the impact resulting from urbanization and industry (Center for Urban Forest Research, 2017). Trees improve air quality, reduce energy consumption, help manage stormwater, reduce erosion, provide critical habitat for wildlife, and promote a connection with nature. When taken together, the community forest contributes to a healthier, more livable, and prosperous Davis.

The City's tree inventory data were analyzed with i-Tree *Eco* benefit-cost modeling software (Eco v6.0.25) to generate the data for this resource analysis. The software uses inventory data collected in the field along with local hourly air pollution and meteorological data to quantify urban forest structure, environmental effects, and value to the community. The program is a central computing engine that makes scientifically sound estimates of the effects of urban forest using peer-reviewed scientific equations to predict environmental and economic benefits. Aesthetic, human health, socio-economic, property value, and wildlife benefits are not calculated as part of this study although they are certainly part of the important benefits provided by Davis' City tree resource.

This report provides an assessment of the structure and composition of the current City tree inventory, consisting of 30,692 trees. Where possible, it also quantifies the benefits derived from the tree resource. This baseline data can be used to make effective resource management decisions, develop policy, and set priorities. Ultimately, the results of the analysis allow the City of Davis to better understand, prioritize, and manage the tree resource.

This summary report provides the following information:

- A description of the current structure of the City tree resource and an established benchmark for future management decisions.
- Quantifiable economic value of benefits from the City tree resource to air quality, stormwater runoff reduction, and carbon sequestration.

- Data that may be used by resource managers in the pursuit of alternative funding sources, local assessment fees, legislative initiatives, and collaborative relationships with utility purveyors, non-governmental organizations, air quality districts, watershed managers, and federal and state agencies.



***Urban trees play an essential role in the community of Davis by providing many benefits, tangible and intangible, to residents, visitors, and neighboring communities.***



# Resource Structure

A tree resource is more thoroughly understood through examination of composition and structure. Consideration of stocking level, species diversity, canopy cover, age distribution, condition, and performance provide a foundation for planning and strategic management. Inferences based on this data can help managers understand the importance of individual trees and species populations to the overall forest as it exists today and provide a basis to plan for and project the future potential of the resource.

## Species Diversity


Species diversity is calculated as the proportion of species representing the total City tree resource (Table 1, Figure 2). The City tree resource includes a mix of 207 unique species (Appendix C), with 17.4% native to California.

**Table 1: Population Summary of Most Prevalent Species**

Species	DBH Class (inches)								# of Trees	% of Pop.
	0 - 4	4 - 8	8 - 12	12 - 18	18 - 24	24 - 30	30 - 36	>36		
<i>Platanus x acerifolia</i>	201	332	550	791	308	124	62	12	2,380	7.75
<i>Pistacia chinensis</i>	266	419	621	530	139	32	8	0	2,016	6.57
<i>Lagerstroemia indica</i>	834	731	207	84	40	13	6	2	1,918	6.25
<i>Quercus lobata</i>	359	306	301	601	161	59	20	25	1,833	5.97
<i>Sequoia sempervirens</i>	59	140	269	635	294	87	21	14	1,518	4.95
<i>Pyrus calleryana</i>	25	138	367	699	205	44	4	2	1,484	4.84
<i>Triadica sebifera</i>	15	99	321	557	235	77	16	3	1,322	4.31
<i>Celtis sinensis</i>	36	105	183	354	206	102	34	8	1,028	3.35
<i>Fraxinus holotricha</i>	12	10	67	323	258	121	25	2	818	2.67
<i>Vitex agnus-castus</i>	64	128	127	144	129	140	50	5	787	2.56
<i>Casuarina cunninghamiana</i>	8	47	62	189	205	103	34	13	661	2.15
<i>Pinus canariensis</i>	7	20	36	139	197	110	47	8	564	1.84
<i>Celtis australis</i>	121	27	48	109	104	76	42	14	541	1.76
<i>Ulmus parvifolia</i>	237	114	77	64	22	10	0	1	525	1.71
<i>Ginkgo biloba</i>	392	36	10	27	33	8	1	0	507	1.65
<i>Acer buergerianum</i>	243	84	52	48	4	0	0	0	431	1.40
<i>Koelreuteria paniculata</i>	26	97	165	100	20	3	1	0	412	1.34
<i>Pinus brutia</i>	32	23	42	69	78	78	42	14	379	1.23
<i>Gleditsia triacanthos</i>	6	26	32	126	130	39	3	1	363	1.18
<i>Quercus douglasii</i>	280	44	22	15	1	0	0	0	362	1.18
<i>Fraxinus velutina</i>	5	3	4	26	91	130	89	6	354	1.15
<i>Quercus suber</i>	71	39	39	69	43	32	19	13	325	1.06
<i>Fraxinus americana</i>	188	54	17	38	18	4	0	1	320	1.04
<i>Platanus racemosa</i>	1	29	59	128	79	11	4	2	313	1.02
All other species	3,152	1,356	1,294	1,854	940	532	241	163	9,531	31.05
<b>Total</b>	<b>6,640</b>	<b>4,407</b>	<b>4,971</b>	<b>7,719</b>	<b>3,941</b>	<b>1,935</b>	<b>769</b>	<b>309</b>	<b>30,692</b>	<b>100%</b>

The species diversity in Davis' City tree resource is higher than the mean of 185 species reported from 18 California communities (Muller and Bornstein, 2010). Four species in the inventory are considered invasive according to California Invasive Species Advisory Committee, including *Triadica sebifera* (Chinese tallowtree), *Schinus molle* (California peppertree), *Ailanthus altissima* (tree of heaven), and *Eucalyptus globulus* (blue gum) (2010).

The most prevalent species are *Platanus x acerifolia* (London planetree, 7.8%), *Pistacia chinensis* (Chinese pistache, 6.6%), *Lagerstroemia indica* (common crapemyrtle, 6.2%), and *Quercus lobata* (California white oak, 6.0%) (Table 1, Figure 2). These four species make up nearly 27% of the overall population. The 24 most prevalent species (representing >1% of the overall population) make up 68.9% of the overall population.



**Figure 2: Species Diversity in Davis' City tree resource**

Maintaining diversity in an urban forest is important. Dominance of any single species or genus can have detrimental consequences in the event of storms, drought, disease, pests, or other stressors that can severely affect a tree resource and the flow of benefits and costs over time. Catastrophic pathogens, such as Dutch elm disease (*Ophiostoma ulmi*), emerald ash borer (*Agilus planipennis*), Asian longhorned beetle (*Anoplophora glabripennis*), and sudden oak death (*Phytophthora ramorum*)



are some examples of unexpected, devastating, and costly pests and pathogens that highlight the importance of diversity and the balanced distribution of species and genera.

Recognizing that all tree species have a potential vulnerability to pests and disease, urban forest managers have long followed a rule of thumb that no single species should represent greater than 10% of the total population and no single genus more than 20% (Santamour, 1990). Among Davis' community tree population, no single species or genera exceed these widely accepted rules. Managers should continue to strive for increased diversity to promote greater resiliency and reduce the risk of a significant loss in benefits should any species become a liability.

## Importance Value

To quantify the significance of any one species in Davis' City tree resource, an importance value (IV) is derived for each of the most prevalent species. Importance values are particularly meaningful to urban forest managers because they indicate a reliance on the functional capacity of a species. **i-Tree Eco calculates importance value based on the sum of two values: percentage of total population and percentage of total leaf area.** Importance value goes beyond tree numbers alone to suggest reliance on specific species based on the benefits they provide. The importance value can range from zero (which implies no reliance) to 200 (suggesting total reliance). A complete table, with importance values for all species, is included in Appendix C.

To reiterate, research strongly suggests that no single species should dominate the composition of an urban forest. Because importance value goes beyond population numbers, it can help managers to better comprehend the resulting loss of benefits from a catastrophic loss of any one species. When importance values are comparatively equal among the 10 to 15 most prevalent species, the risk of significant reductions to benefits is reduced. Of course, suitability of the dominant species is another important consideration. Planting short-lived or poorly adapted species can result in short rotations and increased long-term management costs.

Table 2 lists the importance values of the most prevalent species in Davis' City tree resource. These 24 species represent 68.9% of the overall population and 71.0% of the total leaf area for a combined importance value of 141.3. Of these, Davis relies most heavily on *Platanus x acerifolia* (London planetree, IV=19.1), followed by *Quercus lobata* (California white oak, IV=13.0), and *Sequoia sempervirens* (coast redwood, IV=11.6). Together these three species represent 18.7% of the inventory and have a combined importance value of 43.7 (21.9% of the total). These species contribute significant benefits and a sense of place and are crucial to the inventory and key to sustaining the benefits provided by the City tree resource, as well as preserving the essence of Davis for years to come.

For some species, low importance values are primarily a function of species stature and/or age distribution. Immature or small-stature species frequently have lower importance values than their representation in the inventory might suggest. This is due to their relatively small leaf area and canopy coverage. For example, *Lagerstroemia indica* (common crapemyrtle), which represents 6.2% of the overall resource and 0.8% of overall leaf area, currently has an importance value of 7.0. This species has a large percentage of the population under 8 inches in diameter (81.5%) and the importance value is not likely to increase over time due to its small stature. In contrast, *Quercus douglasii* (blue oak, IV=1.4) represents 1.2% of the resource and less than 1% of overall leaf area. In total, 89.5% of these large stature trees are currently under 8 inches in diameter. As these young trees mature and increase in canopy (leaf area), the importance value of this species is likely to increase significantly over time.

Some species are more significant contributors to the urban forest than population numbers would suggest. For example, *Pinus canariensis* (Canary Island pine) represents 1.8% of the population and 4.0% of overall leaf area and has an importance value of 5.9. This large-stature species is well-established in Davis, with 29.3% of trees greater than 24 inches in diameter. As a result, these trees provide significant benefits despite their representation in the population.

**Table 2: Importance Value (IV) of Prevalent Species in Davis (Representing >1%)**

Species	% of Pop.	% of Leaf Area	Importance Value (IV)
<i>Platanus x acerifolia</i>	7.75	11.30	19.10
<i>Pistacia chinensis</i>	6.57	3.50	10.10
<i>Lagerstroemia indica</i>	6.25	0.80	7.00
<i>Quercus lobata</i>	5.97	7.00	13.00
<i>Sequoia sempervirens</i>	4.95	6.60	11.60
<i>Pyrus calleryana</i>	4.84	3.40	8.20
<i>Triadica sebifera</i>	4.31	3.80	8.10
<i>Celtis sinensis</i>	3.35	6.10	9.40
<i>Fraxinus holotricha</i>	2.67	5.00	7.70
<i>Vitex agnus-castus</i>	2.56	2.40	5.00
<i>Casuarina cunninghamiana</i>	2.15	1.70	3.90
<i>Pinus canariensis</i>	1.84	4.00	5.90
<i>Celtis australis</i>	1.76	3.00	4.80
<i>Ulmus parvifolia</i>	1.71	0.90	2.60
<i>Ginkgo biloba</i>	1.65	0.60	2.30
<i>Acer buergerianum</i>	1.40	0.70	2.10
<i>Koelreuteria paniculata</i>	1.34	1.60	3.00
<i>Pinus brutia</i>	1.23	1.70	2.90
<i>Gleditsia triacanthos</i>	1.18	1.40	2.60
<i>Quercus douglasii</i>	1.18	0.20	1.40
<i>Fraxinus velutina</i>	1.15	1.90	3.10
<i>Quercus suber</i>	1.06	2.00	3.00
<i>Fraxinus americana</i>	1.04	0.50	1.60
<i>Platanus racemosa</i>	1.02	1.90	2.90
All other species	31.05	27.10	58.90
<b>Total</b>	<b>100%</b>	<b>100%</b>	<b>200</b>

## Canopy Cover


The amount and distribution of leaf surface area is the driving force behind the urban forest's ability to produce benefits for the community (Clark et al, 1997). As canopy cover increases, so do the benefits afforded by leaf area. i-Tree Eco estimates that City trees are providing approximately 21.2% canopy cover.

# Relative Age Distribution


Age distribution can be approximated by considering the DBH range of the overall inventory and of individual species. Trees with smaller diameters tend to be younger. It is important to note that palms do not increase in DBH over time and that height more accurately correlates to age.

The distribution of individual tree ages within a tree population influences present and future costs as well as the flow of benefits. An ideally aged population allows managers to allocate annual maintenance costs uniformly over many years and assures continuity in overall tree canopy coverage and associated benefits. A desirable distribution has a high proportion of young trees to offset establishment and age-related mortality as the percentage of older trees declines over time (Richards, 1982/83). This ideal, albeit uneven, distribution suggests a large fraction of trees (~40%) should be young, with a DBH less than eight inches, while only 10% should be in the large diameter classes (>24 inches DBH).

The age distribution of the City tree resource shows a well-established population. Nearly 36% of all trees are less than 8 inches in diameter and 9.8% are greater than 24 inches (Figure 3). The data indicates that a number of recent tree plantings have been directed towards both large and small statured trees.



**Figure 3: Community Tree Inventory Relative Age Distribution**



**Figure 4: Relative Age Distribution of Davis' Top 10 Most Prevalent Species**

Relative age distribution can also be evaluated for each individual species. The 10 most prevalent community tree species are compared against the ideal distribution in Figure 4. Similar to the overall distribution, the majority of the age distributions of the top 10 most prevalent species show well established populations, heavily represented by trees between 12 and 18 inches in diameter (e.g., *Platanus x acerifolia* [London planetree], *Quercus lobata* [California white oak] and *Sequoia sempervirens* [coast redwood]). *Pistacia chinensis* (Chinese pistache) has a nearly ideal age distribution.


The age distribution of *Vitex agnus-castus* (chaste tree) shows a significant portion of the population greater than 24 inches in diameter (24.8%). This species can be a small tree or shrub. The data indicates that most of the population is taking on a shrub form, based on the seemingly large diameter, which is likely a reflection of the method used to measure multi-stemmed individuals (e.g., sum of diameters of individual trunks). *Lagerstroemia indica* (common crapemyrtle) is another small-statured species, therefore trees larger than 8 inches (18.3%) are likely mature. These species do not contribute much to the overall environmental benefits of the tree resource due to their smaller canopies.

Analysis of the age distribution of prevalent species can help resource managers to understand and foresee maintenance activities and budgetary needs. In addition to informing managers of the economics of prevalent species, managers can use the age distribution to determine trends in plantings and adopt strategies for species selection in the years to come.

## Tree Condition & Relative Performance

Tree condition is an indication of how well trees are managed and how well they are performing in the region and in each site-specific environment (e.g., street, median, parking lot, etc.). Condition ratings can help managers anticipate maintenance and funding needs. In addition, tree condition is an important factor for the calculation of resource benefits. A condition rating of good assumes that a tree has no major structural problems, no significant mechanical damage, and may have only minor aesthetic, insect, disease, or structural problems, and is in good health. When trees are performing at their peak, as those rated as good or better, the benefits they provide are maximized.

Based on the inventory data (2018), City trees in Davis are in overall fair or better condition (93.3%). Approximately 5.7% of trees are in poor condition and 1.0% are dead (Figure 5).



**Figure 5: Tree Condition**

### *Relative Performance Index*

The relative performance index (RPI) is one way to further analyze the condition and suitability of a specific tree species. The RPI provides an urban forest manager with a detailed perspective on how different species perform compared to each other. The index compares the condition ratings of each tree species with the condition rating of every other tree species within the inventory. An RPI of 1.0 or better indicates that the species is performing as well or better than average. An RPI value below 1.0 indicates that the species is underperforming in comparison to the rest of the population.

Among Davis' 24 most prevalent tree species, 16 have an RPI of 1.0 or greater (Table 3). *Acer buergerianum* (Trident maple) has the highest RPI at 1.13. The most abundant species, *Platanus x acerifolia* (London planetree, 7.8%) has an RPI of 0.98, which is attributable to 52.2% of the species being in fair condition.

The RPI can be a useful tool for urban forest managers as an indicator of environmental suitability for species selection. If a community has been planting two or more new species, the RPI can be used to compare their relative performance. If the RPI indicates that one is performing relatively poorly, managers may decide to reduce or even stop planting that species and subsequently save money on both planting stock and replacement costs. For example, *Pinus canariensis* (Canary Island pine) has an RPI of 1.07 and *Pinus brutia* (Turkish pine) has an RPI of 0.88 (Table 3). The data indicates that both species are heavily represented by trees between 12 and 18 inches in diameter, and the data indicates that both species have been planted recently. However, the RPI indicates that *P. canariensis* is a more suitable species for Davis where a large-stature coniferous tree is preferred.

The RPI enables managers to look at the performance of long-standing species as well. Established species with an RPI of 1.00 or greater have performed well over time. These top performers should be retained, and planted, as a healthy proportion of the overall population. It is important to keep in mind



that, because RPI is based on condition at the time of the inventory, it may not reflect cosmetic or nuisance issues, especially seasonal issues that are not threatening the health or structure of the trees.

**Table 3: Relative Performance Index of Most Prevalent Species**

Species	Excellent (%)	Good (%)	Fair (%)	Poor (%)	Dead (%)	RPI	# of Trees	% of Pop.
<i>Platanus x acerifolia</i>	0.00	42.60	52.20	4.70	0.40	<b>0.98</b>	2,380	7.75
<i>Pistacia chinensis</i>	0.00	79.30	18.50	1.80	0.30	<b>1.09</b>	2,016	6.57
<i>Lagerstroemia indica</i>	0.00	59.00	36.20	4.60	0.20	<b>1.03</b>	1,918	6.25
<i>Quercus lobata</i>	0.00	46.00	49.40	3.90	0.60	<b>0.99</b>	1,833	5.97
<i>Sequoia sempervirens</i>	0.00	54.20	42.00	2.90	1.00	<b>1.01</b>	1,518	4.95
<i>Pyrus calleryana</i>	0.00	8.40	65.60	24.10	2.00	<b>0.81</b>	1,484	4.84
<i>Triadica sebifera</i>	0.00	54.50	42.10	3.10	0.40	<b>1.02</b>	1,322	4.31
<i>Celtis sinensis</i>	0.20	55.40	38.10	5.50	0.80	<b>1.01</b>	1,028	3.35
<i>Fraxinus holotricha</i>	0.00	31.40	60.10	8.10	0.40	<b>0.94</b>	818	2.67
<i>Vitex agnus-castus</i>	0.00	77.50	19.90	2.50	0.00	<b>1.09</b>	787	2.56
<i>Casuarina cunninghamiana</i>	0.00	71.60	26.80	1.20	0.50	<b>1.07</b>	661	2.15
<i>Pinus canariensis</i>	0.00	73.60	22.30	3.90	0.20	<b>1.07</b>	564	1.84
<i>Celtis australis</i>	0.00	83.20	14.80	1.50	0.60	<b>1.10</b>	541	1.76
<i>Ulmus parvifolia</i>	0.00	78.70	18.90	1.90	0.60	<b>1.09</b>	525	1.71
<i>Ginkgo biloba</i>	0.20	88.60	7.30	1.60	2.40	<b>1.10</b>	507	1.65
<i>Acer buergerianum</i>	0.00	90.30	9.00	0.50	0.20	<b>1.13</b>	431	1.40
<i>Koelreuteria paniculata</i>	0.20	66.50	27.70	5.10	0.50	<b>1.05</b>	412	1.34
<i>Pinus brutia</i>	0.00	12.90	78.40	7.90	0.80	<b>0.88</b>	379	1.23
<i>Gleditsia triacanthos</i>	0.00	19.60	71.10	9.40	0.00	<b>0.90</b>	363	1.18
<i>Quercus douglasii</i>	0.00	77.90	21.00	0.60	0.60	<b>1.09</b>	362	1.18
<i>Fraxinus velutina</i>	0.00	5.60	66.70	27.10	0.60	<b>0.80</b>	354	1.15
<i>Quercus suber</i>	0.00	68.90	26.20	3.70	1.20	<b>1.05</b>	325	1.06
<i>Fraxinus americana</i>	0.00	60.00	31.60	7.50	0.90	<b>1.02</b>	320	1.04
<i>Platanus racemosa</i>	0.00	27.20	66.10	5.80	1.00	<b>0.92</b>	313	1.02
All other species	0.00	0.88	0.84	0.14	0.06	<b>1.02</b>	9,531	31.05
<b>Total</b>	<b>&lt;1.00%</b>	<b>53.10%</b>	<b>40.20%</b>	<b>5.70%</b>	<b>1.00%</b>	<b>1.00</b>	<b>30,692</b>	<b>100%</b>

An RPI value less than 1.00 may be indicative of a species that is not well-adapted to local conditions. Poorly adapted species are more likely to present increased safety and maintenance issues. Species with an RPI less than 1.00 should receive careful consideration before being selected for future planting choices. However, prior to selecting or deselecting trees based on RPI alone, managers should consider the age distribution of the species, among other factors. A species that has an RPI of less than 1.00 but has a significant number of trees in larger DBH classes, may simply be exhibiting signs of population senescence. For example, *Pinus brutia* (Turkish pine), has an RPI of 0.88. This species has a relatively large number of mature trees, with 35.4% larger than 24 inches in diameter. Although the RPI is below 1.00, it is likely an indication of the mature age distribution and small percentage of new plantings

(only 2.5% of trees are below 8 inches in diameter), rather than poor performance. A complete table, with RPI values for all species, is included in Appendix C.

RPI is also helpful for identifying underused species that are demonstrating reliable performance. Species with an RPI value greater than 1.00 and an established age distribution may be indicating their suitability for the local environment. These species should receive consideration for additional planting (Table 4).

As an example, *Celtis australis* (European hackberry) has an RPI of 1.10 and an age distribution that is adequately represented by young to mature trees (27.9% are less than 8 inches in diameter and 24.4% are greater than 24 inches in diameter). The representation in the population and the age distribution combined support the high RPI. Alternatively, *Ulmus davidiana v. japonica* (Japanese elm) represents less than 1% of the population, has an RPI of 1.12, and is almost entirely represented by trees less than 8 inches in diameter (92.6%) and does not have any trees greater than 24 inches in diameter. Although expected to do well in Davis, the current age distribution cannot substantiate the high RPI as there are not enough mature trees, resulting in a lack of evidence for long-term performance.

**Table 4: Species That May Be Underused (based on RPI and age distribution)**

Species	RPI	% of Pop.
<b>Conifer Evergreen Large</b>		
<i>Cedrus deodara</i>	1.04	0.89
<i>Pinus pinea</i>	1.03	0.18
<b>Broadleaf Evergreen Medium</b>		
<i>Eucalyptus nicholii</i>	1.02	0.02
<b>Broadleaf Deciduous Large</b>		
<i>Gymnocladus dioica</i>	1.12	0.57
<i>Quercus macrocarpa</i>	1.13	0.09
<b>Broadleaf Deciduous Medium</b>		
<i>Melia azedarach</i>	1.08	0.41
<b>Broadleaf Deciduous Small</b>		
<i>Cercis canadensis</i>	1.00	0.79

RPI is most relevant when there is a moderately high representation of the species. In other words, if there is a single individual that has a high RPI (greater than 1.00) but is the only representative of the species at the site, additional trial plantings of the species can help test the accuracy of the RPI. It is important to use RPI as one of many factors for species selection. Species that have historically experienced major issues in Davis should be avoided and species with a proven track record should be favored.

## Replacement Value

The replacement value of the existing City tree resource is more than \$91.5 million. Replacement value accounts for the historical investment in trees over their lifetime and is a way of describing the value of a tree population (and/or average value per tree) at a given time. In other words, the value of a tree

is equal to the cost of replacing the tree in its current state (Cullen, 2002). There are several methods available for obtaining a fair and reasonable perception of a tree's value (Council of Tree and Landscape Appraisers, 2018; Watson, 2002). For this analysis, the replacement value reflects current population numbers and is based on the valuation procedures of the Council of Tree and Landscape Appraisers, which uses tree species, diameter, condition, and location information (Nowak et al 2002a; 2002b).

To replace all 30,692 City trees in Davis with trees of equivalent size and condition would cost over \$91.5 million, an average of \$2,983 per tree (Table 5). *Platanus x acerifolia* (London planetree) has the highest replacement value of approximately \$8.1 million and accounts for the greatest proportion of the overall replacement value (8.9%). This is consistent with the species having the highest importance value in the inventory and having well-established age distribution.

The replacement value for Davis' City tree resource reflects the vital importance of these assets to the community. With proper care and maintenance, the value will continue to increase over time. It is important to recognize that replacement values are separate and distinct from the value of annual benefits produced by this resource and in some instances the replacement value of a tree may be greater than or less than the benefits that a particular tree may provide.

**Table 5: Replacement Value for Most Prevalent Species**

Species	Number of Trees	Replacement Value (\$)	% of Replacement Value	% of Pop.
<i>Platanus x acerifolia</i>	2,380	8,143,297	8.90	7.75
<i>Pistacia chinensis</i>	2,016	4,495,144	4.91	6.57
<i>Lagerstroemia indica</i>	1,918	1,880,088	2.05	6.25
<i>Quercus lobata</i>	1,833	5,746,913	6.28	5.97
<i>Sequoia sempervirens</i>	1,518	6,110,424	6.67	4.95
<i>Pyrus calleryana</i>	1,484	3,564,743	3.89	4.84
<i>Triadica sebifera</i>	1,322	4,620,540	5.05	4.31
<i>Celtis sinensis</i>	1,028	4,078,906	4.46	3.35
<i>Fraxinus holotricha</i>	818	3,656,562	3.99	2.67
<i>Vitex agnus-castus</i>	787	3,651,017	3.99	2.56
<i>Casuarina cunninghamiana</i>	661	3,575,834	3.91	2.15
<i>Pinus canariensis</i>	564	3,870,910	4.23	1.84
<i>Celtis australis</i>	541	2,573,654	2.81	1.76
<i>Ulmus parvifolia</i>	525	847,016	0.93	1.71
<i>Ginkgo biloba</i>	507	508,968	0.56	1.65
<i>Acer buergerianum</i>	431	367,193	0.40	1.40
<i>Koelreuteria paniculata</i>	412	822,126	0.90	1.34
<i>Pinus spp.</i>	379	1,721,986	1.88	1.23
<i>Gleditsia triacanthos</i>	363	1,408,791	1.54	1.18
<i>Quercus douglasii</i>	362	156,517	0.17	1.18
<i>Fraxinus velutina</i>	354	2,321,000	2.54	1.15
<i>Quercus suber</i>	325	1,833,490	2.00	1.06
<i>Fraxinus americana</i>	320	421,963	0.46	1.04
<i>Platanus racemosa</i>	313	1,235,475	1.35	1.02
All other species	9,531	23,930,290	26.14	31.05
<b>Total</b>	<b>30,692</b>	<b>\$91,542,848</b>	<b>100%</b>	<b>100%</b>

# Resource Benefits

Community trees continuously mitigate the effects of urbanization and development and protect and enhance the quality of life within the community. The amount and distribution of leaf surface area is the driving force behind the ability of the urban forest to produce benefits for the community (Clark et al, 1997). Healthy trees are vigorous, often producing more leaf surface area each year.

The quantifiable benefits from the urban forest are based on the environmental functions trees perform. In addition to air quality benefits, trees slow down stormwater and remove pollutants, resulting in reduced stormwater management costs for municipalities. Tree growth sequesters carbon in woody stems and roots. The economic value of these ecosystem functions is calculated in terms of both volume and cost savings. It is important to note that this assessment does not fully account for all of the benefits trees provide. For example, i-Tree *Eco* requires information on the distance and aspect of individual trees from homes and other conditioned structures to calculate energy benefits. This information is currently unavailable for Davis' City tree resource. In addition, i-Tree *Eco* does not calculate benefit values for trees larger than 100 inches in diameter. Some trees in the inventory exceeded the maximum allowable diameter and were therefore assigned a default measurement of 100 inches in diameter to accommodate the analysis.

Annual environmental benefits tend to increase with an increase in the number and size of healthy trees (Nowak et al, 2002). Through proper management, urban forest values can be increased over time as trees mature and with improved longevity and as stocking levels are increased. Climate, pest, and weather events can cause values to decrease if the amount of healthy tree cover declines. Excluding energy benefits, the City tree resource provides quantifiable annual environmental benefits valued at approximately \$213,857 (Appendix B).

## Air Quality

Urban trees improve air quality in five fundamental ways:

- Absorption of gaseous pollutants such as ozone (O<sub>3</sub>), sulfur dioxide (SO<sub>2</sub>), and nitrogen dioxide (NO<sub>2</sub>) through leaf surfaces
- Reduction of emissions from power generation by reducing energy consumption
- Increase of oxygen levels through photosynthesis
- Transpiration of water and shade provision, resulting in lower local air temperatures, thereby reducing ozone levels Interception of particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>)<sup>2</sup>




Figure 6: Annual Air Pollution Benefits

<sup>2</sup> PM<sub>2.5</sub> is particulate matter less than 2.5 microns (a subset of PM<sub>10</sub>). These microscopic particles are significant air pollutants and are generally more impactful on human health than PM<sub>10</sub> (i-Tree *Eco* User Manual, 2019).

Air pollutants are known to contribute adversely to human health. Trees decrease the amount of air pollutants in the atmosphere, which can reduce the incidence of numerous negative health effects (Table 6). Ozone is an air pollutant that is particularly harmful to human health. Davis' City trees reduce adverse health effects associated with ozone by approximately 22 incidents annually, a value of \$50,558. Ozone forms when nitrogen oxide from fuel combustion and volatile organic gasses from evaporated petroleum products react in the presence of sunshine. In the absence of cooling effects provided by trees, higher temperatures contribute to ozone formation. In addition to consequences to human health, short-term increases in ozone concentrations are statistically associated with increased tree mortality for 95 large US cities (Bell et al, 2004).

**Table 6: Adverse Health Incidents Avoided Due to Changes in Pollutant Concentration Levels and Economic Values**

	NO <sub>2</sub>		O <sub>3</sub>		PM <sub>2.5</sub>		SO <sub>2</sub>	
	Incidence (Reduction/yr.)	Value (\$/yr.)	Incidence (Reduction/yr.)	Value (\$/yr.)	Incidence (Reduction/yr.)	Value (\$/yr.)	Incidence (Reduction/yr.)	Value (\$/yr.)
Acute Bronchitis					0.00	\$0.37		
Acute Myocardial Infarction					0.00	\$115.88		
Acute Respiratory Symptoms	0.23	\$7.27	17.73	\$1,515.52	3.54	\$346.75	0.02	\$0.56
Asthma Exacerbation	3.45	\$289.29			1.95	\$158.66	0.15	\$11.68
Chronic Bronchitis					0.00	\$542.63		
Emergency Room Visits	0.00	\$0.98	0.00	\$1.83	0.00	\$0.58	0.00	\$0.22
Hospital Admissions	0.01	\$160.41	0.01	\$229.71			0.00	\$14.23
Hospital Admissions, Cardiovascular					0.00	\$19.31		
Hospital Admissions, Respiratory					0.00	\$13.90		
Lower Respiratory Symptoms					0.05	\$2.61		
Mortality			0.01	\$48,353.95	0.01	\$47,754.39		
School Loss Days			4.66	\$457.29				
Upper Respiratory Symptoms					0.04	\$1.82		
Work Loss Days					0.61	\$105.77		
<b>Total</b>	<b>3.69</b>	<b>\$457.95</b>	<b>22.40</b>	<b>\$50,558.30</b>	<b>6.21</b>	<b>\$49,062.67</b>	<b>0.17</b>	<b>\$26.68</b>

**Deposition, Interception, & Avoided Pollutants**

Each year, 1,981.5 pounds of nitrogen dioxide, carbon monoxide, sulfur dioxide, small particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>), and ozone are intercepted or absorbed by City trees, for a total value of \$66,865 (7). As a population, *Platanus x acerifolia* (London planetree) is the greatest contributor to pollutant deposition and interception accounting for 11.3% of the benefit. This is directly related to the species prevalence in the overall population and contributions to the overall leaf area (11.3%).


Trees produce oxygen during photosynthesis, and City trees in Davis produce an estimated 1,124 tons of oxygen annually. Additionally, trees contribute to energy savings by reducing air pollutant emissions (NO<sub>2</sub>, PM<sub>2.5</sub>, SO<sub>2</sub>, and VOCs) that result from energy production.

**Table 7: Annual Air Pollution Removal Benefits**

Air Pollutant	Annual Removal (lb.)	Annual Value (\$)
O <sub>3</sub>	13,947.53	\$50,558
PM <sub>2.5</sub>	267.05	\$49,063
PM <sub>10</sub>	4,993.11	\$16,929
NO <sub>2</sub>	860.47	\$458
CO	538.56	\$388
SO <sub>2</sub>	252.37	\$27
	<b>1,918.45</b>	<b>\$66,865</b>







**Figure 7: Top 5 Species for Air Pollution Removal Benefits**

**Table 8: Annual Air Quality Benefits by Most Prevalent Species**

Species	Number of Trees	% of Pop.	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Platanus x acerifolia</i>	2,380	7.75	1.18	13,327
<i>Pistacia chinensis</i>	2,016	6.57	0.37	4,140
<i>Lagerstroemia indica</i>	1,918	6.25	0.08	99
<i>Quercus lobata</i>	1,833	5.97	0.73	8,233
<i>Sequoia sempervirens</i>	1,518	4.95	0.69	7,801
<i>Pyrus calleryana</i>	1,484	4.84	0.36	3,999
<i>Triadica sebifera</i>	1,322	4.31	0.40	4,510
<i>Celtis sinensis</i>	1,028	3.35	0.63	7,136
<i>Fraxinus holotricha</i>	818	2.67	0.52	5,868
<i>Vitex agnus-castus</i>	787	2.56	0.25	2,832
<i>Casuarina cunninghamiana</i>	661	2.15	0.18	2,002
<i>Pinus canariensis</i>	564	1.84	0.42	4,7212
<i>Celtis australis</i>	541	1.76	0.31	3,530
<i>Ulmus parvifolia</i>	525	1.71	0.10	1,082
<i>Ginkgo biloba</i>	507	1.65	0.07	754
<i>Acer buergerianum</i>	431	1.40	0.08	853
<i>Koelreuteria paniculata</i>	412	1.34	0.17	1,921
<i>Pinus spp.</i>	379	1.23	0.18	1,983
<i>Gleditsia triacanthos</i>	363	1.18	0.15	1,636
<i>Quercus douglasii</i>	362	1.18	0.02	249
<i>Fraxinus velutina</i>	354	1.15	0.20	2,253
<i>Quercus suber</i>	325	1.06	0.21	2,338
<i>Fraxinus americana</i>	320	1.04	0.05	597
<i>Platanus racemosa</i>	313	1.02	0.19	2,178
All other species	9,531	31.05	2.82	32,571
<b>Total</b>	<b>30,692</b>	<b>100%</b>	<b>10.43</b>	<b>\$117,423</b>

While trees do a great deal to absorb air pollutants (especially ozone and particulate matter); they also negatively contribute to air pollution. Trees emit volatile organic compounds (VOCs), which also contribute to ozone and carbon monoxide formation. The i-Tree *Eco* analysis accounts for these VOC emissions in the air quality cumulative benefit. Trees in Davis are estimated to emit 27,594 pounds of volatile organic compounds (VOCs) (15,401 pounds of isoprene and 12,192 pounds of monoterpenes) annually. Emissions vary based on species characteristics (e.g., some genera such as oaks are high isoprene emitters) and amount of leaf biomass. The highest volume of VOC emissions is generated by *Quercus lobata* (California white oak), accounting for approximately 25.1% of the overall emissions, largely due to their size (7.0% of overall leaf area) and species attributes. Regardless, the net air quality benefit of *Quercus lobata* is positive.

Air quality impacts of trees are complex, and the i-Tree *Eco* software models these interactions to help urban forest managers evaluate the true impact of urban trees on the Davis' air quality. The cumulative and interactive effects of trees on climate, pollution removal, VOCs, and power plant emissions determine the net impact of trees on air pollution. Local urban forest management decisions also can help improve air quality by prioritizing tree species recognized for their ability to improve air quality and planting next to large traffic corridors.

## Atmospheric Carbon Dioxide Reductions


As environmental awareness continues to increase, conversations around global warming and the effects of greenhouse gas (GHG) emissions are increasing. As energy from the sun (sunlight) strikes the Earth's surface it is reflected into space as infrared radiation (heat). GHGs absorb some of this infrared radiation and trap heat in the atmosphere, modifying the temperature of the Earth's surface. Many chemical compounds in the Earth's atmosphere act as GHGs, including carbon dioxide (CO<sub>2</sub>), water vapor, and human-made (gases/aerosols). As GHGs increase, the amount of energy radiated back into space is reduced, and more heat is trapped in the atmosphere. An increase in the average temperature of the Earth may result in changes in weather, sea levels, and land-use patterns, commonly referred to as "climate change" (NASA, 2020).

Because urban trees use carbon as a building component for wood and foliar growth, they can help offset carbon emissions and should be recognized as a part of a community's solution for meeting carbon offset goals identified in climate action plans and other environmental policies. i-Tree tools can be used to estimate the GHG and carbon sequestration benefits of tree planting projects (California Air Resource Board, 2020).

Urban trees reduce atmospheric CO<sub>2</sub> in two ways:

- Directly, through growth and the sequestration of CO<sub>2</sub> in wood, foliar biomass, and soil
- Indirectly, by lowering the demand for heating and air conditioning, thereby reducing the emissions associated with electric power generation and natural gas consumption

To date, Davis' City trees are estimated to have stored 16,158 tons of carbon (CO<sub>2</sub>) in woody and foliar biomass valued at nearly \$2.8 million. Annually, the City tree resource directly sequesters an additional 421.5 tons of carbon valued at \$71,882, with an average value of \$2.00 per tree (Table 9). Among prevalent species, *Casuarina cunninghamiana* (river she-oak, \$7.87/tree), *Triadica sebifera* (Chinese tallowtree, \$5.24/tree), and *Gleditsia triacanthos* (honeylocust, \$3.93/tree) provide the greatest annual per-tree benefits to atmospheric carbon removal, sequestering 13,562.3 tons of carbon annually (Figure 8). These three species account for 18.9% of overall carbon benefit and 7.6% of the overall population.



**Figure 8: Top 5 Species for Carbon Benefits**

**Table 9: Annual Carbon Sequestration Benefits by Most Prevalent Species**

Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Average \$/tree	% of Annual Benefit
<i>Platanus x acerifolia</i>	2,380	7.75	51.27	8,744.03	3.67	12.16
<i>Pistacia chinensis</i>	2,016	6.57	16.54	2,820.84	1.40	3.92
<i>Lagerstroemia indica</i>	1,918	6.25	9.87	1,682.83	0.88	2.34
<i>Quercus lobata</i>	1,833	5.97	28.02	4,778.93	2.61	6.65
<i>Sequoia sempervirens</i>	1,518	4.95	21.13	3,603.16	2.37	5.01
<i>Pyrus calleryana</i>	1,484	4.84	24.73	4,217.58	2.84	5.87
<i>Triadica sebifera</i>	1,322	4.31	40.65	6,932.12	5.24	9.64
<i>Celtis sinensis</i>	1,028	3.35	3.28	559.13	0.54	0.78
<i>Fraxinus holotracha</i>	818	2.67	18.35	3,129.21	3.83	4.35
<i>Vitex agnus-castus</i>	787	2.56	4.52	771.33	0.98	1.07
<i>Casuarina cunninghamiana</i>	661	2.15	30.51	5,203.75	7.87	7.24
<i>Pinus canariensis</i>	564	1.84	9.80	1,671.14	2.96	2.32
<i>Celtis australis</i>	541	1.76	1.63	277.64	0.51	0.39
<i>Ulmus parvifolia</i>	525	1.71	7.61	1,298.44	2.47	1.81
<i>Ginkgo biloba</i>	507	1.65	0.89	152.03	0.30	0.21
<i>Acer buergerianum</i>	431	1.40	2.72	464.13	1.08	0.65
<i>Koelreuteria paniculata</i>	412	1.34	4.95	844.62	2.05	1.18
<i>Pinus spp.</i>	379	1.23	6.17	1,051.66	2.77	1.46
<i>Gleditsia triacanthos</i>	363	1.18	8.36	1,426.45	3.93	1.98
<i>Quercus douglasii</i>	362	1.18	0.65	110.83	0.31	0.15
<i>Fraxinus velutina</i>	354	1.15	5.42	923.86	2.61	1.29
<i>Quercus suber</i>	325	1.06	4.43	755.23	2.32	1.05
<i>Fraxinus americana</i>	320	1.04	2.26	385.45	1.20	0.54
<i>Platanus racemosa</i>	313	1.02	2.62	447.07	1.43	0.62
All other species	9,531	31.05	115.08	19,630.56	1.95	27.31
<b>Total</b>	<b>30,692</b>	<b>100%</b>	<b>421.46</b>	<b>\$71,882.02</b>	<b>\$2.00</b>	<b>100%</b>

## Stormwater Runoff Reductions


Rainfall interception by trees reduces the amount of stormwater that enters collection and treatment facilities during large storm events (Figure 9). Trees intercept rainfall in their canopy, acting as mini reservoirs, controlling runoff at the source. Healthy urban trees reduce the amount of runoff and pollutant loading in receiving waters in three primary ways:

- Leaves and branch surfaces intercept and store rainfall, thereby reducing runoff volumes and delaying the onset of peak flows
- Root growth and decomposition increase the capacity and rate of soil infiltration by rainfall and reduce overland flow
- Tree canopies reduce soil erosion and surface flows by diminishing the impact of raindrops on bare soil

Davis' City tree resource is estimated to contribute to the avoidance of more than 189 million gallons of stormwater runoff annually through the interception of precipitation on the leaves and bark of trees for an average of 6,160 gallons per tree (Table 10). The total value of this benefit is \$24,552 annually, an average of \$0.80 per tree.

*Platanus x acerifolia* (London planetree) provide 11.4% of the estimated total avoided runoff and provide the greatest per tree benefit of \$1.17 (Table 10, Figure 10). Their age distribution and stature allow them to provide a larger benefit in comparison to other species. In contrast, *Lagerstroemia indica* (common crapemyrtle), which represents 3.6% of the population, reduce less than 1% of the estimated total avoided runoff. This small stature species is limited in its ability to intercept stormwater. Characteristics that contribute to greater stormwater capture include large leaves, broad or dense canopies, and furrowed bark.

As trees grow, the benefits that they provide tend to grow as well. Some species provide more benefits than others, based on their architecture and leaf morphology. Other trees have characteristics that hinder their ability to be strong contributors to stormwater runoff reduction, possibly due to a tree having smaller leaves and thinner canopies.



**Figure 9: How Trees Impact Stormwater**




Figure 10: Top 5 Species for Stormwater Benefits

Table 10: Stormwater Benefits by Most Prevalent Tree Species

Species Name	Number of Trees	Leaf Area (acres)	Potential Evapotranspiration (gallons/yr.)	Evaporation (gallons/yr.)	Transpiration (gallons/yr.)	Water Intercepted (gallons/yr.)	Avoided Runoff (gallons/yr.)	Avoided Runoff Value (\$)
<i>Platanus x acerifolia</i>	2,380	154	21,459,727	1,528,125	9,130,095	1,530,345	311,843	2,787
<i>Pistacia chinensis</i>	2,016	48	6,666,415	474,709	2,836,243	475,398	96,873	866
<i>Lagerstroemia indica</i>	1,918	10	1,462,912	104,172	622,400	104,324	21,258	190
<i>Quercus lobata</i>	1,833	95	13,256,693	943,996	5,640,094	945,367	192,640	1,721
<i>Sequoia sempervirens</i>	1,518	90	12,561,410	894,485	5,344,284	895,784	182,537	1,631
<i>Pyrus calleryana</i>	1,484	46	6,438,664	458,491	2,739,346	459,157	93,564	836
<i>Triadica sebifera</i>	1,322	52	7,262,083	517,126	3,089,671	517,877	105,529	943
<i>Celtis sinensis</i>	1,028	82	11,491,029	818,265	4,888,888	819,453	166,982	1,492
<i>Fraxinus holotricha</i>	818	68	9,448,929	672,849	4,020,071	673,826	137,307	1,227
<i>Vitex agnus-castus</i>	787	33	4,559,529	324,679	1,939,863	325,151	66,257	592
<i>Casuarina cunninghamiana</i>	661	23	3,224,194	229,592	1,371,742	229,925	46,853	419
<i>Pinus canariensis</i>	564	54	7,603,084	541,408	3,234,752	542,194	110,485	987
<i>Celtis australis</i>	541	41	5,684,398	404,780	2,418,442	405,368	82,603	738
<i>Ulmus parvifolia</i>	525	12	1,741,671	124,023	740,998	124,203	25,309	226
<i>Ginkgo biloba</i>	507	9	1,213,733	86,429	516,386	86,554	17,637	158
<i>Acer buergerianum</i>	431	10	1,374,084	97,847	584,607	97,989	19,968	178
<i>Koelreuteria paniculata</i>	412	22	3,092,627	220,223	1,315,766	220,543	44,941	402
<i>Pinus brutia</i>	379	23	3,193,385	227,398	1,358,634	227,728	46,405	415
<i>Gleditsia triacanthos</i>	363	19	2,633,956	187,561	1,120,623	187,834	38,275	342
<i>Quercus douglasii</i>	362	3	400,882	28,546	170,556	28,588	5,825	52
<i>Fraxinus velutina</i>	354	26	3,628,393	258,374	1,543,709	258,750	52,726	471
<i>Quercus suber</i>	325	27	3,763,907	268,024	1,601,364	268,413	54,695	489
<i>Fraxinus americana</i>	320	7	960,973	68,430	408,848	68,529	13,964	125
<i>Platanus racemosa</i>	313	25	3,507,740	249,783	1,492,377	250,145	50,973	455
All other species	9,531	376	52,446,214	3,734,641	22,313,374	3,740,066	762,125	6,810
<b>Total</b>	<b>30,692</b>	<b>1,355</b>	<b>189,076,631</b>	<b>13,463,956</b>	<b>80,443,132</b>	<b>13,483,511</b>	<b>2,747,575</b>	<b>\$24,552.30</b>





## Energy Savings

Trees modify climate and conserve energy in three principal ways:

- Shading reduces the amount of radiant energy absorbed and stored by hardscape surfaces, thereby reducing the heat island effect
- Transpiration converts moisture to water vapor, thereby cooling the air by using solar energy that would otherwise result in heating of the air
- Reduction of wind speed plus the movement of outside air into interior spaces, and conductive heat loss where thermal conductivity is relatively high (e.g., glass windows) (Simpson, 1998)

The heat island effect describes the increase in urban temperatures in relation to surrounding suburban and rural areas. Heat islands are associated with an increase in hardscape and impervious surfaces. Trees and other vegetation within an urbanized environment help reduce the heat island effect by lowering air temperatures 5°F (3°C) compared with outside the green space (Chandler, 1965). On a larger scale, temperature differences of more than 9°F (5°C) have been observed between city centers without adequate canopy coverage and more vegetated suburban areas (Akbari et al, 1997). The relative importance of these effects depends upon the size and configuration of trees and other landscape elements (McPherson, 1993). Tree spacing, crown spread, and vertical distribution of leaf area each influence the transport of warm air and pollutants along streets and out of urban canyons. Trees reduce conductive heat loss from buildings by reducing air movement into buildings and against conductive surfaces (e.g., glass, metal siding). Trees can reduce wind speed and the resulting air infiltration by up to 50%, translating into potential annual heating savings of 25% (Heisler, 1986).

### *Electricity & Natural Gas Reductions*

Energy reduction metrics are calculated using data on tree distance and direction from buildings. The annual energy reductions from Davis' City trees were not calculated because this data is not currently captured in the inventory database. However, trees in Davis contribute to electric and natural gas savings through shading and climate buffering effects.

## Aesthetic, Property Value, & Socioeconomic Benefits

Trees provide beauty in the urban landscape, privacy and screening, improved human health, a sense of comfort and place, and habitat for urban wildlife. Research shows that trees promote better business by stimulating more frequent and extended shopping and a willingness to pay more for goods and parking (Wolf, 2007). In residential areas, the values of these benefits are captured as a percentage of the value of the property on which a tree stands. There is no current model for calculating the aesthetic benefits of an urban forest. Although, there are many indicators that suggest trees and tree canopy cover contribute significantly to quality of life and community well-being.

It is important to acknowledge that this assessment does not account for all the benefits provided by the tree resource. Some benefits are intangible and/or difficult to quantify, such as:

- Impacts on psychological and physical health and wellness
- Reduction in crime and violence
- Increases in tourism revenue
- Quality of life
- Wildlife habitat

- Socio-economic impacts
- Increases in property values
- Overall community well-being

Empirical evidence of these benefits does exist (Wolf, 2007; Kaplan and Kaplan, 1989; Ulrich, 1986), but there is limited knowledge about the physical processes at work and the complex nature of interactions make quantification imprecise. Tree growth and mortality rates are highly variable. A true and full accounting of benefits and investments must consider variability among sites (e.g., tree species, growing conditions, maintenance practices), as well as variability in tree growth. In other words, trees are worth far more than what one can ever quantify!

## Calculating Tree Benefits

While all these tree benefits are provided by the community forest, it can be useful to understand the contribution of just one tree. Individuals can calculate the benefits of individual trees to their property by using i-Tree *Design* ([design.itreetools.org](http://design.itreetools.org)).




## Annual Benefits of Most Prevalent Species

It is important to keep in mind that a benefits analysis provides a snapshot of the City tree inventory as it exists today. The calculated benefits are based on the size and condition of existing trees. To provide greater context for the overall per tree and per species benefits of the most prevalent tree species (Figure 11, Table 11), and to determine if these benefits are a true indicator of performance, the age distribution and stature of the species must also be considered (Table 1, Figure 4).

The most prevalent tree species in Davis' City tree resource, *Platanus x acerifolia* (London planetree, 7.8%) is providing the greatest overall annual benefit, a value of \$24,858, which is attributable to its prevalence in the population as well as species characteristics (Figure 11). The data indicates recent plantings of this large-statured, well-established species in the inventory, which is important because as this population ages, maintenance needs (and costs) may increase and per tree benefits will begin to level out.

Among other prevalent species, *Pinus canariensis* (Canary Island pine) provides \$7,380 in annual benefits and the highest per tree benefit, an average of \$13.09 per tree. In contrast, *Quercus douglasii* (blue oak) provides \$411.85 in annual benefits and the lowest per tree benefit, an average of \$1.14 per tree. As the majority (89.5%) of *Quercus douglasii* measure less than 8 inches in diameter, which for this large-statured species are not yet mature, the benefits that this large-statured species provides are likely to increase significantly over time as trees grow and mature.



**Figure 11: Summary of Annual Benefits for Most Prevalent Species**



**Table 11: Summary of Annual Benefits of Most Prevalent Species**


Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Avoided Runoff (gallon/yr.)	Avoided Runoff (\$/yr.)	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Platanus x acerifolia</i>	2,380	7.75	51.27	8,744	311,843	2,786.63	1.18	13,327
<i>Pistacia chinensis</i>	2,016	6.57	16.54	2,821	96,873	865.66	0.37	4,140
<i>Lagerstroemia indica</i>	1,918	6.25	9.87	1,683	21,258	189.96	0.08	909
<i>Quercus lobata</i>	1,833	5.97	28.02	4,779	192,640	1,721.43	0.73	8,233
<i>Sequoia sempervirens</i>	1,518	4.95	21.13	3,603	182,537	1,631.15	0.69	7,801
<i>Pyrus calleryana</i>	1,484	4.84	24.73	4,218	93,564	836.08	0.36	3,999
<i>Triadica sebifera</i>	1,322	4.31	40.65	6,932	105,529	943.01	0.40	4,510
<i>Celtis sinensis</i>	1,028	3.35	3.28	559	166,982	1,492.15	0.63	7,136
<i>Fraxinus holotricha</i>	818	2.67	18.35	3,129	137,307	1,226.98	0.52	5,868
<i>Vitex agnus-castus</i>	787	2.56	4.52	771	66,257	592.07	0.25	2,832
<i>Casuarina cunninghamiana</i>	661	2.15	30.51	5,204	46,853	418.67	0.18	2,002
<i>Pinus canariensis</i>	564	1.84	9.80	1,671	110,485	987.29	0.42	4,722
<i>Celtis australis</i>	541	1.76	1.63	278	82,603	738.14	0.31	3,530
<i>Ulmus parvifolia</i>	525	1.71	7.61	1,298	25,309	226.16	0.10	1,082
<i>Ginkgo biloba</i>	507	1.65	0.89	152	17,637	157.61	0.07	754
<i>Acer buergerianum</i>	431	1.40	2.72	464	19,968	178.43	0.08	853
<i>Koelreuteria paniculata</i>	412	1.34	4.95	845	44,941	401.59	0.17	1,921
<i>Pinus spp.</i>	379	1.23	6.17	1,052	46,405	414.67	0.18	1,983
<i>Gleditsia triacanthos</i>	363	1.18	8.36	1,426	38,275	342.03	0.15	1,636
<i>Quercus douglasii</i>	362	1.18	0.65	111	5,825	52.06	0.02	249
<i>Fraxinus velutina</i>	354	1.15	5.42	924	52,726	471.16	0.20	2,253
<i>Quercus suber</i>	325	1.06	4.43	755	54,695	488.76	0.21	2,338
<i>Fraxinus americana</i>	320	1.04	2.26	385	13,964	124.79	0.05	597
<i>Platanus racemosa</i>	313	1.02	2.62	447	50,973	455.49	0.19	2,178
All other species	9,531	31.05	115.08	19,631	762,125	6,810.33	2.82	32,571
<b>Total</b>	<b>30,692</b>	<b>100%</b>	<b>421.47</b>	<b>\$71,882</b>	<b>2,747,575</b>	<b>\$24,552.30</b>	<b>10.43</b>	<b>\$117,423</b>

# Net Annual Benefits

Davis receives substantial benefits from their City tree resource; however, managers should understand and evaluate the investment required to preserve the City tree resource along with the benefits that it provides. A limitation of the annual benefits summary is that i-Tree *Eco* does not fully account for all benefits provided by City tree resource. Many of the documented environmental and socioeconomic benefits provided by trees are intangible and not able to be quantified using current methods (University of Washington, 2018; University of Illinois, 2018).

Davis' City tree resource has a beneficial effect on the environment, and annually contributes \$213,857 in quantifiable benefits to the community (Figure 12). Individual components of the environmental benefits include improved air quality \$117,423 (54.9%), carbon reduction of \$71,882 (33.6%), and stormwater management for \$24,552 (11.5%).

Annually, City trees provide a total benefit of \$213,857, a value of \$6.97 per tree and \$3.09 per capita.



### Annual Investment & Benefit Offset

Figure 12: Annual Environmental Benefits

Davis' urban forestry staff provided estimated investment costs. The total annual cost of managing the City tree resource in Davis is approximately \$1.6 million. Based on budget information from 2021 and 2022, in total, 40.8% of the costs are attributed to annual pruning, 32.9% to tree removal, and 8.0% to purchasing and planting trees. The remaining 18% of costs are for program administration and claims. The quantifiable benefits from i-Tree *Eco* offset this investment by \$213,857 (Table 12).

Table 12: Quantifiable Benefits and Investments

Benefits	Total (\$)	\$/tree	\$/capita
Carbon Sequestration	71,882	2.34	1.04
Pollution Removal	117,423	3.83	1.69
Avoided Runoff	24,552	0.80	0.35
<b>Total Benefits</b>	<b>\$213,857</b>	<b>\$6.97</b>	<b>\$3.09</b>

Investments	Total (\$)	\$/tree	\$/capita
Purchasing Trees and Planting	127,061	4.14	1.83
Contract Pruning	643,235	20.96	9.28
Removal	519,604	16.93	7.50
Administration	255,711	8.33	3.69
Liability/Claims	32,284	1.05	0.47
<b>Total Investments</b>	<b>\$1,577,895</b>	<b>\$51.41</b>	<b>\$22.77</b>

## Pest and Pathogen Threats

Management of pests and disease organisms can be a challenge in any urban forest. In some cases, a pest or disease can result in significant tree damage or loss and/or be costly to manage. Involvement in the global economy, close proximity to major ports, and a highly mobile human population increase the risk of an invasive pest or pathogen introduction into Davis. To further investigate the risk of pests and pathogens, i-Tree *Eco* identifies the susceptibility of tree populations to 36 emerging and existing pests and pathogens in the United States (Appendix B). According to the analysis, 21,162 (69.0%) of Davis' City trees are susceptible to the included pests and pathogens and the potential risk is estimated at nearly \$65.0 million. The pests and pathogens identified as most relevant to Davis are included in Table 13. Anticipating and monitoring for these threats is an important part of urban forest management.

Among the pests of greatest concern for Davis' community forest is the polyphagous shot hole borer. The polyphagous shot hole borer is involved in a disease called Fusarium dieback, which occurs when invasive beetles feed on fungi that they carry into heartwood tissues of the tree. Some of the introduced fungi are tree pathogens that disrupt the flow of water and nutrients. Staining and gummosis can be seen around beetle entry and exit wounds, and typically cankers have formed at these sites. The damage causes branch dieback, and over time can kill the tree (Eskalen, 2018). Within the United States, the polyphagous shot hole borer has been detected in southern California but has the potential to spread to the Central Valley as these beetles have a large host range consisting of more than 260 plant species and can colonize healthy or stressed trees. An estimated 46.5% of trees in Davis are at risk to polyphagous shot hole borer.

Defoliating moths, such as spongy moth (*Lymantria dispar*) and winter moth (*Operophtera brumata*), are not yet present in California, but they threaten a broad range of tree hosts present in Davis (22.3% and 15.2% of trees susceptible, respectively). During outbreaks, the feeding damage weakens the tree host, and renders it more vulnerable to other pests and diseases (Collins, 1996). The gypsy moth is known to feed on hundreds of species of trees and shrubs; oaks (*Quercus*) are one of their preferred hosts.

Davis is currently experiencing thousand cankers disease (TCD) in the walnut populations and Dutch elm disease in the elm trees populations. In TCD, the walnut twig beetle (*Pityophthorus juglandis*) vectors the fungus *Geosmithia morbida*. As the walnut twig beetle tunnels into the cambium, cankers form in and around beetle galleries. When the beetles are abundant, cankers can girdle twigs or branches, stopping the flow of sugars through the phloem, and causing yellowing, wilting, and branch die back (Tisserat et al, 2009). Trees under stress usually die within three years of initial symptoms but sometimes the infection can persist for years with no external signs or symptoms. Control measures have not yet been identified, but sanitation practices to dispose of infected material is advised.

Dutch Elm Disease (DED) has devastated *Ulmus americana* (American elm) populations, which are some of the most important street trees in the twentieth century. Since first reported in the 1930s, it has killed over 50% of the native elm population in the United States (Forest Service, Northeastern Area State and Private Forestry, 2005). Less than 1% of Davis' City tree inventory is susceptible to DED, largely because managers have planted elm species that exhibit some disease resistance.

**Table 13: Pest & Pathogen Threats to Davis**

		Number of Trees		Structural Value (\$)		Leaf Area (%)		Leaf Area (ac)	
		Susceptible	Not Susceptible	Susceptible	Not Susceptible	Susceptible	Not Susceptible	Susceptible	Not Susceptible
aspen leafminer	<i>Phyllocnistis populiella</i>	162	30,530	306,416	91,236,432	0.30	99.70	4.30	1,350.50
Asian longhorned beetle	<i>Anoplophora glabripennis</i>	2,661	28,031	3,307,601	88,235,247	4.50	95.50	60.90	1,293.90
butternut canker	<i>Ophiognomonia clavignenti-juglandacearum</i>	14	30,678	105,619	91,437,229	0.10	99.90	1.20	1,353.50
Dutch elm disease	<i>Ophiostoma ulmi</i>	177	30,515	303,734	91,239,114	0.30	99.70	4.30	1,350.50
emerald ash borer	<i>Agrilus planipennis</i>	2,022	28,670	8,396,883	83,145,965	9.90	90.10	133.50	1,221.20
fusiform rust	<i>Cronartium quercuum</i>	23	30,669	50,330	91,492,518	0.10	99.90	0.90	1,353.90
goldspotted oak borer	<i>Agrilus auroguttatus</i>	244	30,448	837,411	90,705,437	1.20	98.80	15.90	1,338.90
large aspen tortrix	<i>Choristoneura conflictana</i>	209	30,483	375,582	91,167,266	0.40	99.60	5.80	1,349.00
laurel wilt	<i>Xyleborus glabratus and Raffaellea lauricola</i>	42	30,650	138,529	91,404,319	0.20	99.80	2.00	1,352.70
mountain pine beetle	<i>Dendroctonus ponderosae</i>	15	30,677	61,313	91,481,535	0.10	99.90	0.80	1,354.00
oak wilt	<i>Ceratocystis fagacearum</i>	4,202	26,490	11,829,292	79,713,556	14.30	85.70	194.40	1,160.40
pine shoot beetle	<i>Tomicus piniperda</i>	1,293	29,399	8,121,462	83,421,386	7.60	92.40	103.20	1,251.60
polyphagous shot hole borer	<i>Euwallacea sp. and Fusarium euwallaceae</i>	14,258	16,434	41,726,417	49,816,431	43.60	56.40	590.50	764.30
spongy moth	<i>Lymantria dispar</i>	6,841	23,851	17,433,314	74,109,534	20.00	80.00	270.60	1,084.20
spruce beetle	<i>Dendroctonus rufipennis</i>	4	30,688	5,369	91,537,479	0.00	100.00	0.10	1,354.70
spruce budworm	<i>Choristoneura fumiferana</i>	4	30,688	5,369	91,537,479	0.00	100.00	0.10	1,354.70
sudden oak death	<i>Phytophthora ramorum</i>	2,414	28,278	8,554,718	82,988,130	9.90	90.10	133.70	1,221.10
southern pine beetle	<i>Dendroctonus frontalis</i>	1,297	29,395	8,126,831	83,416,017	7.60	92.40	103.20	1,251.60
sirex wood wasp	<i>Sirex noctilio</i>	1,293	29,399	8,121,462	83,421,386	7.60	92.40	103.20	1,251.60
thousand cankers disease	<i>Geosmithia morbida and Pityophthorus juglandis</i>	215	30,477	1,140,480	90,402,368	1.10	98.90	15.00	1,339.80
winter moth	<i>Operophtera brumata</i>	4,679	26,013	12,317,111	79,225,737	14.40	85.60	194.50	1,160.30
western pine beetle	<i>Dendroctonus brevicomi</i>	2	30,690	4,380	91,538,468	0.00	100.00	0.00	1,354.70
western spruce budworm	<i>Choristoneura occidentalis</i>	18	30,674	63,432	91,479,416	0.10	99.90	0.80	1,354.00
<b>All Pests</b>		<b>21,162</b>	<b>9,530</b>	<b>\$64,987,241</b>	<b>\$26,555,607</b>	<b>69.50</b>	<b>30.50</b>	<b>941.40</b>	<b>413.40</b>





## Pest Management

Although managers cannot foresee when a pest or pathogen may be introduced to the urban forest, being aware of potential threats is the first step in a preparedness program. Following Integrated Pest Management (IPM) protocol and best management practices when preparing for and addressing pest and diseases can help to minimize their economic, health, and environmental consequences (Wiseman and Raupp, 2016). Some management practices include:

- Obtain current information on emergent pests and pathogens
- Increase understanding of the biology of the pest and pathogen as well as the tree symptoms that indicate infestation/infection
- Identify procedures and protocols that will be followed in the case of an introduced pest or pathogen
- Complete training and licensing in the case of pesticide or fungicide use
- Plant tree species that are resistant or tolerant to identified pest and pathogen threats
- Choose healthy, vigorous nursery stock
- Diversify plantings at the genus level, as many pests threaten several species within a genus
- Prevent the movement of felled tree materials that may be harboring pests or pathogens such as untreated logs, firewood, and woodchips



***Maintaining a diverse City tree resource is important in integrated pest management.***

## Conclusion

This analysis describes the current structural characteristics of Davis' City tree resource, using established numerical modeling and statistical methods to provide a general accounting of the benefits. The analysis provides a "snapshot" of this resource at its current population, structure, and condition. Trees are providing quantifiable impacts on air quality, reduction in atmospheric CO<sub>2</sub>, stormwater runoff, and aesthetic benefits. Davis' 30,692 City trees provide cumulative annual benefits worth \$213,857, a value of \$667 per tree and \$3.09 per capita.

Industry standards suggest that no one tree species should represent more than 10% of the urban forest. According to the 2018 inventory data, no species in Davis violate this rule. Additionally, industry standards suggest no one genera should represent more than 20% of a population. Of Davis' City tree inventory, no genus violates this rule. The rule provides a baseline for greater genetic diversity, therefore future new and replacement tree plantings should continue to focus on increasing the diversity of the City tree resource.

Davis' City tree resource has an established population in fair or better condition with 207 distinct species. The City should continue to focus resources on preserving existing and mature trees to promote health, strong structure, and tree longevity. Structural and training pruning for young trees will maximize the value of this resource, reduce long-term maintenance costs, reduce risk, and ensure that as trees mature, they provide the greatest possible benefits over time.

Based on this resource analysis, DRG recommends the following regarding the management of the City's trees:


- Protect existing trees and regularly inspect trees to identify and mitigate structural and age-related defects.
- Provide structural pruning for young trees and continue to work toward a routine pruning cycle for all trees.
- Monitor species performance (e.g., health, structure, longevity, pest and disease resistance) and increase resilience in the urban forest by planting species that perform best in local and regional conditions, including introducing new species that indicate promising traits.
  - Consider species performance data when reviewing and updating the Climate Ready Tree List.
  - Discourage the planting of species that are classified as invasive (e.g., *Triadica sebifera* [Chinese tallowtree], *Schinus molle* [California peppertree], *Ailanthus altissima* [tree of heaven], and *Eucalyptus globulus* [blue gum]).
- Maintain the benefits of key species by continuing to include them in new tree plantings.
- Increase genus and species diversity in new and replacement tree plantings to reduce reliance on over-represented species. While no species represent more than 10% of the overall population, increasing diversity in the tree resource can provide additional benefits.
- Prioritize planting replacement trees for those trees that are removed.
- Consider successional planting of important species, as supported by relative performance index (RPI) and the relative age distribution (e.g., *Celtis sinensis* [Chinese hackberry]).
- Use new tree plantings to improve diversity, increase benefits, and support an ideal age distribution of community trees.



- Plant large-stature species for greater benefits wherever space allows.
- Follow best management practices when monitoring for and dealing with pests and diseases.
- Maintain and update the inventory database to include new tree plantings, removals, as well as changes in diameter, condition for new trees.
  - Consider adding information on distance and orientation to nearest structure/building so that energy benefits can be calculated in future analyses.
  - Consider adding locations for available planting sites.

Urban forest managers can better anticipate future trends with an understanding of the composition and structure of the tree population. Managers can also anticipate challenges and devise plans to increase benefits. Performance data from this analysis can be used to make determinations regarding species selection, distribution, and maintenance policies. Documenting current structure is necessary for establishing goals and performance objectives and can serve as a benchmark for measuring future success.

Davis' City trees are of vital importance to the environmental, social, and economic well-being of the community. Inventory data can be used to plan a proactive and forward-looking approach to the care of City trees. Updates should continue to be incorporated into the inventory as regular maintenance is performed, including information on the diameter and condition of existing trees. Current and complete inventory data will help staff to track maintenance activities and tree health more efficiently and will provide a strong basis for making informed management decisions. A continued commitment to planting, maintaining, and preserving these trees will support the health and welfare of the City and the community at large.



***Trees are of vital importance to the environmental, social, and economic well-being of the community.***

## Appendix A: References

- Akbari, H., D. Kurn, et al. 1997. Peak power and cooling energy savings of shade trees. *Energy and Buildings* 25:139–148.
- British Columbia Ministry of Water, Land, and Air Protection. 2005. Residential wood burning emissions in British Columbia. British Columbia Carbon Dioxide Information Analysis Center. 2010. <https://cdiac.ess-dive.lbl.gov/home.html>
- California Air Resource Board. 2020. Urban and Community Forestry program Quantification Methodology. Retrieved from: [https://ww2.arb.ca.gov/sites/default/files/classic/cc/capandtrade/auctionproceeds/calfire\\_ucf\\_finalq\\_m\\_012820.pdf](https://ww2.arb.ca.gov/sites/default/files/classic/cc/capandtrade/auctionproceeds/calfire_ucf_finalq_m_012820.pdf)
- California Invasive Species Advisory Committee. 2010. The California Invasive Species List. CA: Invasive Species Council of California. <<http://www.iscc.ca.gov/docs/CaliforniaInvasiveSpeciesList.pdf>>
- Center for Urban Forest Research. 2017. Retrieved from [https://www.fs.fed.us/psw/topics/urban\\_forestry/](https://www.fs.fed.us/psw/topics/urban_forestry/)
- Chandler T.J. 1965. *The Climate of London*. London UK. Hutchinson.
- Clark JR, Matheny NP, Cross G, Wake V. 1997. A model of urban forest sustainability. *Journal of Arboriculture* 23 (1): 17-30.
- Collins, J. 1996. European Gypsy Moth. University of Kentucky Entomology Fact Sheet-425. Lexington, KY. Retrieved from <https://entomology.ca.uky.edu/ef425>
- Council of Tree and Landscape Appraisers. 2018. *Guide for Plant Appraisal*. 10th Edition. International Society of Arboriculture.
- Cullen S. 2002. Tree Appraisal: Can Depreciation Factors Be Rated Greater than 100%? *Journal of Arboriculture* 28(3):153-158.
- [EIA] Energy Information Administration. 2001. Total Energy Consumption in U.S. Households by Type of Housing Unit. <http://www.eia.doe.gov/emeu/recs/contents.html>.
- [EIA] Environmental Protection Agency (EPA), 2015; Interagency Working Group on Social Cost of Carbon. [www3.epa.gov/climatechange/Downloads/EPAactivities/social-cost-carbon.pdf](http://www3.epa.gov/climatechange/Downloads/EPAactivities/social-cost-carbon.pdf)
- EPA. 2015. Social Cost of Carbon. [https://www.epa.gov/sites/production/files/201612/documents/social\\_cost\\_of\\_carbon\\_fact\\_sheet.pdf](https://www.epa.gov/sites/production/files/201612/documents/social_cost_of_carbon_fact_sheet.pdf)
- Eskalen, A. 2018. Identifying Polyphagous and Kuroshio Shot Hole Borer in California. <https://anrcatalog.ucanr.edu/pdf/8590.pdf>
- Georgia Forestry Commission. 2009. Biomass Energy Conversion for Electricity and Pellets Worksheet. Dry Branch, GA: Georgia Forestry Commission
- Graham, R.L., Wright, L.L., and Turhollow, A.F. 1992. The potential for short-rotation woody crops to reduce U.S. CO<sub>2</sub> Emissions. *Climatic Change* 22:223-238.
- Heisler, G.M. 1986. Energy Savings with Trees. *J Arbor* 12 (5): 113–125.
- Kaplan R., and Kaplan S. 1989. *The Experience of Nature: A Psychological Perspective*. Cambridge: Cambridge University Press.



Layton, M. 2004. 2005 Electricity Environmental Performance Report: Electricity Generation and Air Emissions. California Energy Commission.

[http://www.energy.ca.gov/2005\\_energypolicy/documents/2004-11-15\\_workshop/2004-11-15\\_03-A\\_LAYTON.PDF](http://www.energy.ca.gov/2005_energypolicy/documents/2004-11-15_workshop/2004-11-15_03-A_LAYTON.PDF)

Leonardo Academy Inc. 2010. Leonardo Academy's Guide to Calculating Emissions Including Emission Factors and Energy Prices. Retrieved from:

<http://www.cleanerandgreener.org/download/Leonardo%20Academy%20C&G%20Emission%20Factors%20and%20Energy%20Prices.pdf>

McPherson E.G. 1993. Evaluating the Cost-Effectiveness of Shade Trees for Demand-Side Management. *Electricity Journal* 6(9):57-65.

McPherson, E.G. and J. R. Simpson 1999. Carbon dioxide reduction through urban forestry: guidelines for professional and volunteer tree planters. Gen. Tech. Rep. PSW-171. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research station 237 p.

[http://wcufr.ucdavis.edu/products/cufr\\_43.pdf](http://wcufr.ucdavis.edu/products/cufr_43.pdf)

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Xiao, Q. 1999. Tree Guidelines for San Joaquin Valley Communities. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Center for Urban Forest Research.

McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Crowell, A.M.N.; Xiao, Q. 2010. Northern California coast community tree guide: benefits, costs, and strategic planting. PSW-GTR-228. Gen. Tech. Rep. PSW-GTR-228. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.

McPherson, E.G., Xiao, X., Maco, S.E., Van Der Zanden, A., Simpson, J.R., Bell, N., Peper, P.J. 2002. Western Washington and Oregon Community Tree Guide: Benefits, Costs and Strategic Planting. Center for Urban Forest Research Pacific Southwest Research Station. [Fs.fed.us/psw](http://Fs.fed.us/psw)

Muller, R.N., and C. Bornstein. 2010. Maintaining the diversity of California's municipal forests *Journal of Arboriculture* 36.1: 18.

NASA, 2020. What is the greenhouse effect? Earth Science Communications Team at NASA's Jet Propulsion Laboratory and the California Institute of Technology. Retrieved from

<https://climate.nasa.gov/faq/19/what-is-the-greenhouse-effect/>

Nowak, D.J. 1995. Trees pollute? A "TREE" explains it all. In: Proceedings of the 7th National Urban Forestry Conference. Washington, DC: American Forests. Pp. 28-30

Nowak, D.J., and D.E. Crane. 2000. The Urban Forest Effects (UFORE) Model: quantifying urban forest structure and functions. In: Hansen, M., and T. Burk (Eds.) Integrated Tools for Natural Resources Inventories in the 21st Century. Proc. Of the IUFRO Conference. USDA Forest Service General Technical Report NC-212. North Central Research Station, St. Paul, MN. Pp. 714-720. See also

<http://www.ufore.org>.

Nowak, D.J.; Crane, D.E.; Dwyer, J.F. 2002a. Compensatory value of urban trees in the United States. *Journal of Arboriculture*. 28(4): 194 – 199.

Nowak, D.J.; Crane, D.E.; Stevens, J.C.; Ibarra, M. 2002b. Brooklyn's Urban Forest. Gen. Tech. Rep. NE-290. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Research Station. 107 p. Council of Tree and Landscape Appraisers guidelines. For more information, see

- Nowak, D.J., D.E. Crane, and J.F. Dwyer. 2002. Compensatory value of urban trees in the United States. *J. Arboric.* 28(4): 194-199.
- Pajaro Valley Arts. Discover Davis. N.d. Retrieved from <https://pvarts.org/discover-Davis/>
- Peper, P.J.; McPherson, E.G.; Simpson, J.R.; Vargas, K.E.; Xiao, Q. 2008. City of Indianapolis, Indiana, Municipal Forest Resource Analysis. Internal Tech. Rep. Davis, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Center for Urban Forest Research. 68 p.
- Richards, N.A. 1982/83. Diversity and Stability in a Street Tree Population. *Urban ecology.* 7:159-171.
- Simpson JR. 1998. Urban Forest Impacts on Regional Space Condition Energy Use: Sacramento County Case Study. *Journal of Arboriculture* 24(4): 201-214.
- Sperling's Best Places, n.d. Davis, California. Retrieved from <https://www.bestplaces.net/climate/city/california/davis>
- Tisserat, N., Cranshaw, W., Leatherman, D., Utley, C., & Alexander, K. (2009). Black walnut mortality in Colorado caused by the walnut twig beetle and thousand cankers disease. *Plant Health Progress*, 10(1), 10.
- Ulrich, R.S. 1986. Human Responses to Vegetation and Landscapes. *Landscape and Urban Planning*, 13, 29-44.
- University of Illinois. 2018. Landscape and Human Health Laboratory. Retrieved from: <http://lhhl.illinois.edu/research.htm>
- University of Washington. 2018. Green Cities: Good Health. Retrieved from: <http://depts.washington.edu/hhwb/>
- US Census Bureau. n.d. QuickFacts Davis, California. Retrieved from <https://www.census.gov/quickfacts/Daviscitycalifornia>
- Vargas, K.E.; McPherson, E.G.; Simpson, J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007a. Interior West community tree guide: benefits, costs, and strategic planting. Gen. Tech. Rep. PSW-GTR-205. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 105 p.
- Vargas, K.E.; McPherson, E.G.; Simpson J.R.; Peper, P.J.; Gardner, S.L.; Xiao, Q. 2007b. Temperate interior West community tree guide: benefits, costs, and strategic planting. Gen. Tech. Rep. PSW-GTR-206. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 108 p.
- Watson, G. 2002. Comparing formula methods of tree appraisal. *Journal of Arboriculture*, 28(1):11-18.
- Wolf, K.L. 2007. The environmental Psychology of Trees. *International Council of Shopping Centers Research Review.* 14, 3:39-43.
- Worrall, J.J. 2007. Chestnut Blight. *Forest and Shade Tree Pathology.* [http://www.forestpathology.org/dis\\_chestnut.html](http://www.forestpathology.org/dis_chestnut.html)



*Davis' City tree resource includes a mix of 207 distinct species.*



## Appendix B: Methods

### i-Tree *Eco* Model and Field Measurements

All field data was collected during the leaf-on season to properly assess tree canopies. The i-Tree *Eco* model uses inventory data, local hourly air pollution, and meteorological data to quantify the urban forest and its structure and benefits (Nowak & Crane, 2000), including:

- Urban forest structure (e.g., genus composition, tree health, leaf area, etc.).
- Amount of pollution removed hourly by the urban forest, and its associated percent air quality improvement throughout a year. Pollution removal is calculated for ozone, sulfur dioxide, nitrogen dioxide, carbon monoxide and particulate matter (<2.5 microns).
- Total carbon stored and net carbon annually sequestered by the urban forest.
- Structural value of the forest as a replacement cost.
- Potential impact of infestations by pests or pathogen.

### Definitions and Calculations

**Avoided surface water runoff** value is calculated based on rainfall interception by vegetation, specifically the difference between annual runoff with and without vegetation. Although tree leaves, branches, and bark may intercept precipitation and thus mitigate surface runoff, only the precipitation intercepted by leaves is accounted for in this analysis. The U.S. value of avoided runoff, \$0.067 per ft<sup>3</sup>, is based on the U.S. Forest Service's Community Tree Guide Series (McPherson et al, 1999-2010; Peper et al, 2009; 2010; Vargas et al, 2007a-2008).

**Carbon dioxide emissions** from automobile assumed six pounds of carbon per gallon of gasoline if energy costs of refinement and transportation are included (Graham et al, 1992).

**Carbon emissions** were calculated based on the total city carbon emissions from the 2010 US per capita carbon emissions (Carbon Dioxide Information Analysis Center, 2010) This value was multiplied by the population of Davis (69,295) to estimate total city carbon emissions.

**Carbon sequestration** is removal of carbon from the air by plants. Carbon storage and carbon sequestration values are calculated based on \$170.55 per short ton (EPA, 2015; Interagency Working Group on Social Cost of Carbon, 2015).

**Carbon storage** is the amount of carbon bound up in the above-ground and below-ground parts of woody vegetation. Carbon storage and carbon sequestration values are calculated based on \$170.55 per ton (EPA, 2015; Interagency Working Group on Social Cost of Carbon, 2015).

**Diameter at Breast Height (DBH)** is the diameter of the tree measured 4'5" above grade.

**Household emissions** average is based on average electricity kWh usage, natural gas Btu usage, fuel oil Btu usage, kerosene Btu usage, LPG Btu usage, and wood Btu usage per household in 2009 (EIA, 2013; EIA, 2014), CO<sub>2</sub>, SO<sub>2</sub>, and NO<sub>3</sub> power plant emission per kWh (Leonardo Academy, 2011), CO emission per kWh assumes 1/3 of one percent of C emissions is CO (EIA, 2014), PM<sub>10</sub> emission per kWh (Layton 2004), CO<sub>2</sub>, NO<sub>3</sub>, SO<sub>2</sub>, and CO emission per Btu for natural gas, propane and butane (average used to represent LPG), Fuel #4 and #6 (average used to represent fuel oil and kerosene) (Leonardo Academy, 2011), CO<sub>2</sub> emissions per Btu of wood (EIA, 2014), CO, NO<sub>3</sub> and SO<sub>2</sub> emission per Btu based

on total emissions and wood burning (tons) from (British Columbia Ministry, 2005; Georgia Forestry Commission, 2009).

**Leaf area** was estimated using measurements of crown dimensions and percentage of crown canopy missing.

**Monetary values (\$)** are reported in US dollars throughout the report.

**Ozone (O<sub>3</sub>)** is an air pollutant that is harmful to human health. Ozone forms when nitrogen oxide from fuel combustion and volatile organic gases from evaporated petroleum products react in the presence of sunshine. In the absence of cooling effects provided by trees, higher temperatures contribute to ozone (O<sub>3</sub>) formation.

**Passenger automobile emissions** assumed 0.72 pounds of carbon per driven mile (U.S. Environmental Protection Agency, 2010) multiplied by the average miles driven per vehicle in 2011 (Federal Highway Administration, 2013).

**Pollution removal** is calculated based on the prices of \$1,442.52 per ton (carbon monoxide), \$7,249.79 per ton (ozone), \$1,064.41 per ton (nitrogen dioxide), \$211.43 per ton (sulfur dioxide), \$367,444.76 per ton (particulate matter less than 2.5 microns), and \$6,780.92 per ton (particulate matter less than 10 microns) (Nowak et al., 2014).

**Potential pest impacts** were estimated based on tree inventory information from the study area combined with i-Tree *Eco* pest range maps. The input data included species, DBH, total height, height to crown base, crown width, percent canopy missing, and crown dieback. In the model, potential pest risk is based on pest range maps and the known pest host species that are likely to experience mortality.

**Pest range maps** for 2012 from the Forest Health Technology Enterprise Team (FHTET) (Forest Health Technology Enterprise Team, 2014) were used to determine the proximity of each pest to Yolo County. For the county, it was established whether the insect/disease occurs within the county, is within 250 miles of the county edge, is between 250 and 750 miles away, or is greater than 750 miles away. FHTET did not have pest range maps for Dutch elm disease and chestnut blight. The range of these pests was based on known occurrence and the host range, respectively (Eastern Forest Environmental Threat Assessment Center; Worrall 2007). Due to the dates of some of these resources, pests may have encroached closer to the tree resource in recent years.

**Replacement value** is based on the physical resource itself (e.g., the cost of having to replace a tree with a similar tree). Structural values were based on valuation procedures of the Council of Tree and Landscape Appraisers, which uses tree species, diameter, condition, and location information (Nowak et al 2002a; 2002b).

**Ton** is equivalent to a U.S. short ton, or 2,000 pounds.



## Appendix C: Tables

**Table 14: Botanical and Common Names of Tree Species in Davis' City tree resource**

Botanical Name	Common Name	# of Trees	% of Pop.
<i>Platanus x acerifolia</i>	London planetree	2,380	7.75
<i>Pistacia chinensis</i>	Chinese pistache	2,016	6.57
<i>Lagerstroemia indica</i>	common crapemyrtle	1,918	6.25
<i>Quercus lobata</i>	California white oak	1,833	5.97
<i>Sequoia sempervirens</i>	coast redwood	1,518	4.95
<i>Pyrus calleryana</i>	Callery pear	1,484	4.84
<i>Triadica sebifera</i>	Chinese tallowtree	1,322	4.31
<i>Celtis sinensis</i>	Chinese hackberry	1,028	3.35
<i>Fraxinus holotricha</i>	Moraine ash	818	2.67
<i>Vitex agnus-castus</i>	chaste tree	787	2.56
<i>Casuarina cunninghamiana</i>	river she-oak	661	2.15
<i>Pinus canariensis</i>	Canary Island pine	564	1.84
<i>Celtis australis</i>	European hackberry	541	1.76
<i>Ulmus parvifolia</i>	Chinese elm	525	1.71
<i>Ginkgo biloba</i>	ginkgo	507	1.65
<i>Acer buergerianum</i>	Trident maple	431	1.40
<i>Koelreuteria paniculata</i>	goldenrain tree	412	1.34
<i>Pinus brutia</i>	Turkish pine	379	1.23
<i>Gleditsia triacanthos</i>	honeylocust	363	1.18
<i>Quercus douglasii</i>	blue oak	362	1.18
<i>Fraxinus velutina</i>	velvet ash	354	1.15
<i>Quercus suber</i>	cork oak	325	1.06
<i>Fraxinus americana</i>	white ash	320	1.04
<i>Platanus racemosa</i>	California sycamore	313	1.02
<i>Koelreuteria bipinnata</i>	Chinese flame tree	305	0.99
<i>Quercus virginiana</i>	live oak	278	0.91
<i>Cedrus deodara</i>	deodar cedar	274	0.89
<i>Acer rubrum</i>	red maple	258	0.84
<i>Cercis canadensis</i>	eastern redbud	244	0.79
<i>Quercus agrifolia</i>	coastal live oak	236	0.77
<i>Ulmus davidiana v. japonica</i>	Japanese elm	232	0.76
<i>Olea europaea</i>	olive	226	0.74
<i>Celtis occidentalis</i>	northern hackberry	222	0.72
<i>Fraxinus uhdei</i>	Shamel ash	214	0.70
<i>Zelkova serrata</i>	Japanese zelkova	212	0.69
<i>Juglans hindsii</i>	hind walnut	201	0.65
<i>Laurus nobilis</i>	bay laurel	198	0.65
<i>Malus spp.</i>	apple species	192	0.63
<i>Quercus wislizeni</i>	interior live oak	189	0.62
<i>Prunus cerasifera</i>	cherry plum	188	0.61

Botanical Name	Common Name	# of Trees	% of Pop.
<i>Arbutus unedo</i>	strawberry tree	187	0.61
<i>Liriodendron tulipifera</i>	tulip tree	187	0.61
<i>Quercus rubra</i>	northern red oak	183	0.60
<i>Eucalyptus sideroxylon</i>	Mugga ironbark	178	0.58
<i>Acer x freemanii</i>	Freeman maple	177	0.58
<i>Gymnocladus dioica</i>	Kentucky coffee tree	176	0.57
<i>Morus alba</i>	white mulberry	172	0.56
<i>Fraxinus angustifolia ssp. Oxycarpa</i>	Caucasian ash	169	0.55
<i>Magnolia grandiflora</i>	southern magnolia	160	0.52
<i>Pinus halepensis</i>	Aleppo pine	157	0.51
<i>Quercus buckleyi</i>	Buckley oak	153	0.50
<i>Quercus ilex</i>	holly oak	152	0.50
<i>Pistacia spp.</i>	pistache species	152	0.50
<i>Ulmus spp.</i>	elm species	145	0.47
<i>Quercus canbyi</i>	Chisos oak	143	0.47
<i>Melia azedarach</i>	Chinaberry	125	0.41
<i>Fraxinus spp.</i>	ash species	124	0.40
<i>Prunus spp.</i>	plum species	122	0.40
<i>x Chitalpa tashkentensis</i>	chitalpa	113	0.37
<i>Tilia cordata</i>	littleleaf linden	95	0.31
<i>Sophora japonica</i>	necklacepod	90	0.29
<i>Quercus shumardii</i>	Shumard oak	86	0.28
<i>Populus fremontii</i>	Fremont cottonwood	85	0.28
<i>Acer platanooides</i>	Norway maple	84	0.27
<i>Pinus spp.</i>	pine species	82	0.27
<i>Robinia pseudoacacia</i>	black locust	82	0.27
<i>Quercus coccinea</i>	scarlet oak	80	0.26
<i>Ligustrum lucidum</i>	glossy privet	78	0.25
<i>Chilopsis linearis</i>	desert-willow	77	0.25
<i>Quercus robur</i>	English oak	77	0.25
<i>Prunus serrulata</i>	Japanese flowering cherry	73	0.24
<i>Searsia lancea</i>	African sumac	66	0.22
<i>Carpinus betulus</i>	European hornbeam	64	0.21
<i>Juglans regia</i>	English walnut	63	0.21
<i>Cupressus arizonica</i>	Arizona cypress	59	0.19
<i>Maclura pomifera</i>	Osage orange	56	0.18
<i>Pinus pinea</i>	Italian stone pine	55	0.18
<i>Populus nigra v. italica</i>	Lombardy poplar	52	0.17
<i>Ostrya virginiana</i>	eastern hophornbeam	52	0.17
<i>Chionanthus retusus</i>	Chinese fringe tree	44	0.14
<i>Schinus molle</i>	California peppertree	44	0.14
<i>Acer campestre</i>	hedge maple	43	0.14
<i>Acacia melanoxylon</i>	blackwood	41	0.13

Botanical Name	Common Name	# of Trees	% of Pop.
<i>Salix spp.</i>	willow species	40	0.13
<i>Pinus nigra</i>	Austrian pine	38	0.12
<i>Eucalyptus camaldulensis</i>	red gum eucalyptus	36	0.12
<i>Liquidambar styraciflua</i>	sweetgum	33	0.11
<i>Calocedrus decurrens</i>	incense cedar	31	0.10
<i>Acer saccharinum</i>	silver maple	31	0.10
<i>Aesculus californica</i>	California buckeye	31	0.10
<i>Albizia julibrissin</i>	Persian silk tree	30	0.10
<i>Quercus macrocarpa</i>	bur oak	29	0.09
<i>Chamaecyparis</i>	false cypress spp.	29	0.09
<i>Geijera parviflora</i>	Australian willow	27	0.09
<i>Robinia spp.</i>	robinia species	26	0.08
<i>Umbellularia californica</i>	California laurel	26	0.08
<i>Chilopsis spp.</i>	desert willow species	25	0.08
<i>Platanus occidentalis</i>	American sycamore	24	0.08
<i>Fraxinus ornus</i>	flowering ash	23	0.07
<i>Prunus dulcis</i>	sweet almond	23	0.07
<i>Quercus spp.</i>	oak species	23	0.07
<i>Tilia tomentosa</i>	silver linden	22	0.07
<i>Quercus frainetto</i>	Hungarian oak	21	0.07
<i>Betula pendula</i>	European white birch	21	0.07
<i>Pinus thunbergii</i>	Japanese black pine	20	0.07
<i>Casuarina stricta</i>	she-oak	19	0.06
<i>Parrotia persica</i>	Persian ironwood	19	0.06
<i>Tilia americana</i>	American basswood	18	0.06
<i>Ulmus americana</i>	American elm	18	0.06
<i>Eucalyptus polyanthemos</i>	silver dollar eucalyptus	17	0.06
<i>Ficus spp.</i>	fig species	17	0.06
<i>Platanus orientalis</i>	Oriental planetree	17	0.06
<i>Cinnamomum camphora</i>	camphor tree	16	0.05
<i>Taxodium distichum</i>	bald cypress	16	0.05
<i>Prunus persica</i>	peach	15	0.05
<i>Tilia x euclora</i>	Crimean linden	15	0.05
<i>Prosopis chilensis</i>	Chilean mesquite	15	0.05
<i>Catalpa speciosa</i>	northern catalpa	15	0.05
<i>Ceratonia siliqua</i>	carob	14	0.05
<i>Juglans nigra</i>	black walnut	14	0.05
<i>Acer palmatum</i>	Japanese maple	14	0.05
<i>Maytenus boaria</i>	mayten	13	0.04
<i>Pinus sylvestris</i>	Scots pine	13	0.04
<i>Paulownia tomentosa</i>	royal paulownia	12	0.04
<i>Tristaniopsis spp.</i>	tristaniopsis species	12	0.04
<i>Quercus cerris</i>	European turkey oak	11	0.04

Botanical Name	Common Name	# of Trees	% of Pop.
<i>Acer negundo</i>	boxelder	11	0.04
<i>Alnus cordata</i>	Italian alder	11	0.04
<i>Cedrus atlantica</i>	atlas cedar	11	0.04
<i>Afrocarpus gracilior</i>	fern pine	11	0.04
<i>Quercus palustris</i>	pin oak	10	0.03
<i>Alnus rhombifolia</i>	white alder	10	0.03
<i>Eriobotrya japonica</i>	loquat tree	10	0.03
<i>Acer truncatum x platanoides</i>	sunset maple	9	0.03
<i>Eucalyptus melliodora</i>	yellow box	9	0.03
<i>Prunus Kanzan</i>	Kanzan cherry	9	0.03
<i>Pterocarya stenoptera</i>	Chinese wingnut	9	0.03
<i>Punica spp.</i>	pomegranate species	9	0.03
<i>Quercus laurifolia</i>	laurel oak	9	0.03
<i>Ulmus procera</i>	English elm	9	0.03
<i>Magnolia x soulangeana</i>	saucer magnolia	8	0.03
<i>Melaleuca linariifolia</i>	cajeput tree	8	0.03
<i>Crataegus spp.</i>	hawthorn species	7	0.02
<i>Eucalyptus spp.</i>	gum species	7	0.02
<i>Ailanthus altissima</i>	tree of heaven	7	0.02
<i>Quercus chrysolepis</i>	canyon live oak	7	0.02
<i>Acer pseudoplatanus</i>	sycamore maple	6	0.02
<i>Eucalyptus nicholii</i>	willow-leaved gimlet	6	0.02
<i>Quercus muehlenbergii</i>	chinkapin oak	6	0.02
<i>Salix babylonica</i>	Babylon weeping willow	6	0.02
<i>Betula nigra</i>	river birch	5	0.02
<i>Carpinus caroliniana</i>	American hornbeam	5	0.02
<i>Quercus imbricaria</i>	shingle oak	5	0.02
<i>Xylosma spp.</i>	xylosma species	5	0.02
<i>Picea pungens</i>	blue spruce	4	0.01
<i>Cupressus sempervirens</i>	Italian cypress	4	0.01
<i>Thuja plicata</i>	western red cedar	4	0.01
<i>Ulmus pumila</i>	Siberian elm	4	0.01
<i>Magnoliopsida</i>	Unknown species	4	0.01
<i>Lagerstroemia subcostata</i>	Chinese crapemyrtle	4	0.01
<i>Myrtuss spp.</i>	myrtus species	4	0.01
<i>Parkinsonia aculeata</i>	Jerusalem thorn	3	0.01
<i>Metasequoia glyptostroboides</i>	dawn redwood	3	0.01
<i>Juglans spp.</i>	walnut species	3	0.01
<i>Eucalyptus globulus</i>	blue gum eucalyptus	3	0.01
<i>Brachychiton populneus</i>	kurrajong	3	0.01
<i>Acer griseum</i>	paperbark maple	3	0.01
<i>Pinus radiata</i>	Monterey pine	3	0.01
<i>Quercus castaneifolia</i>	chestnut-leaved oak	2	0.01

Botanical Name	Common Name	# of Trees	% of Pop.
<i>Photinia spp.</i>	chokeberry species	2	0.01
<i>Quercus nigra</i>	water oak	2	0.01
<i>Washingtonia filifera</i>	California palm	2	0.01
<i>Prunus x yedoensis</i>	Yoshino flowering cherry	2	0.01
<i>Prunus lyonii</i>	Catalina cherry	2	0.01
<i>Prunus armeniaca</i>	apricot	2	0.01
<i>Pinus coulteri</i>	Coulter pine	2	0.01
<i>Crataegus phaenopyrum</i>	Washington hawthorn	2	0.01
<i>Corymbia maculata</i>	spotted gum	2	0.01
<i>Citrus spp.</i>	citrus species	2	0.01
<i>Citrus limon</i>	lemon	2	0.01
<i>Callistemon spp.</i>	bottlebrush sprvird	2	0.01
<i>Acer saccharum</i>	Sugar maple	2	0.01
<i>Acer spp.</i>	maple species	2	0.01
<i>Quercus phellos</i>	willow oak	2	0.01
<i>Koelreuteria elegans</i>	flamegold	2	0.01
<i>Thuja spp.</i>	red cedar species	1	0.00
<i>Persea spp.</i>	bay species	1	0.00
<i>Pistacia vera</i>	pistachio	1	0.00
<i>Prunus subhirtella</i>	Higan cherry	1	0.00
<i>Prunus x blireiana</i>	Blireiana plum	1	0.00
<i>Pyrus betulifolia</i>	birchleaf pear	1	0.00
<i>Ostrya carpinifolia</i>	Hop hornbeam	1	0.00
<i>Tetradium spp.</i>	tetradium species	1	0.00
<i>Firmiana simplex</i>	Chinese parasoltree	1	0.00
<i>Ulmus x hollandica</i>	Dutch elm	1	0.00
<i>Quercus engelmannii</i>	Engelmann oak	1	0.00
<i>Nyssa sylvatica</i>	black tupelo	1	0.00
<i>Magnolia virginiana</i>	sweetbay	1	0.00
<i>Malus x arnoldiana</i>	Arnold's apple	1	0.00
<i>Juniperus virginiana</i>	eastern red cedar	1	0.00
<i>Elaeocarpus decipiens</i>	Japanese blueberry tree	1	0.00
<i>Cercidiphyllum japonicum</i>	katsura tree	1	0.00
<i>Carya illinoensis</i>	pecan	1	0.00
<i>Brachychiton acerifolius</i>	Illwarra flame tree	1	0.00
<i>Acer macrophyllum</i>	bigleaf maple	1	0.00
<i>Washingtonia spp.</i>	palm species	1	0.00
<i>Larix occidentalis</i>	western larch	1	0.00
<b>Total</b>		<b>30,692</b>	<b>100%</b>



**Table 15: Population Summary for All Species**

Species	DBH Class (inches)								# of Trees	% of Pop.
	0 - 4	4 - 8	8 - 12	12 - 18	18 - 24	24 - 30	30 - 36	>36		
<i>Platanus x acerifolia</i>	201	332	550	791	308	124	62	12	2,380	7.75
<i>Pistacia chinensis</i>	266	419	621	530	139	32	8	0	2,016	6.57
<i>Lagerstroemia indica</i>	834	731	207	84	40	13	6	2	1,918	6.25
<i>Quercus lobata</i>	359	306	301	601	161	59	20	25	1,833	5.97
<i>Sequoia sempervirens</i>	59	140	269	635	294	87	21	14	1,518	4.95
<i>Pyrus calleryana</i>	25	138	367	699	205	44	4	2	1,484	4.84
<i>Triadica sebifera</i>	15	99	321	557	235	77	16	3	1,322	4.31
<i>Celtis sinensis</i>	36	105	183	354	206	102	34	8	1,028	3.35
<i>Fraxinus holotricha</i>	12	10	67	323	258	121	25	2	818	2.67
<i>Vitex agnus-castus</i>	64	128	127	144	129	140	50	5	787	2.56
<i>Casuarina cunninghamiana</i>	8	47	62	189	205	103	34	13	661	2.15
<i>Pinus canariensis</i>	7	20	36	139	197	110	47	8	564	1.84
<i>Celtis australis</i>	121	27	48	109	104	76	42	14	541	1.76
<i>Ulmus parvifolia</i>	237	114	77	64	22	10	0	1	525	1.71
<i>Ginkgo biloba</i>	392	36	10	27	33	8	1	0	507	1.65
<i>Acer buergerianum</i>	243	84	52	48	4	0	0	0	431	1.40
<i>Koelreuteria paniculata</i>	26	97	165	100	20	3	1	0	412	1.34
<i>Pinus brutia</i>	32	23	42	69	78	78	42	14	379	1.23
<i>Gleditsia triacanthos</i>	6	26	32	126	130	39	3	1	363	1.18
<i>Quercus douglasii</i>	280	44	22	15	1	0	0	0	362	1.18
<i>Fraxinus velutina</i>	5	3	4	26	91	130	89	6	354	1.15
<i>Quercus suber</i>	71	39	39	69	43	32	19	13	325	1.06
<i>Fraxinus americana</i>	188	54	17	38	18	4	0	1	320	1.04
<i>Platanus racemosa</i>	1	29	59	128	79	11	4	2	313	1.02
<i>Koelreuteria bipinnata</i>	243	15	10	21	12	4	0	0	305	0.99
<i>Quercus virginiana</i>	33	38	40	118	46	2	0	1	278	0.91
<i>Cedrus deodara</i>	32	17	31	99	46	33	12	4	274	0.89
<i>Acer rubrum</i>	213	29	11	4	1	0	0	0	258	0.84
<i>Cercis canadensis</i>	146	28	24	19	10	16	1	0	244	0.79
<i>Quercus agrifolia</i>	45	39	39	78	22	8	4	1	236	0.77
<i>Ulmus davidiana v japonica</i>	182	33	13	3	1	0	0	0	232	0.76
<i>Olea europaea</i>	20	32	27	67	43	19	13	5	226	0.74
<i>Celtis occidentalis</i>	41	48	40	65	19	5	1	3	222	0.72
<i>Fraxinus uhdei</i>	10	11	16	67	60	32	15	3	214	0.70
<i>Zelkova serrata</i>	156	28	10	8	3	7	0	0	212	0.69
<i>Juglans hindsii</i>	6	20	16	32	26	29	22	50	201	0.65
<i>Laurus nobilis</i>	25	73	48	36	5	4	3	4	198	0.65
<i>Malus spp.</i>	77	62	45	7	0	1	0	0	192	0.63
<i>Quercus wislizeni</i>	13	45	33	58	24	13	3	0	189	0.62
<i>Prunus cerasifera</i>	27	36	106	17	0	0	1	1	188	0.61
<i>Arbutus unedo</i>	33	66	56	17	11	4	0	0	187	0.61
<i>Liriodendron tulipifera</i>	11	32	70	66	7	1	0	0	187	0.61
<i>Quercus rubra</i>	160	10	12	1	0	0	0	0	183	0.60

Species	DBH Class (inches)								# of Trees	% of Pop.
	0 - 4	4 - 8	8 - 12	12 - 18	18 - 24	24 - 30	30 - 36	>36		
<i>Eucalyptus sideroxylon</i>	1	2	15	68	60	23	8	1	178	0.58
<i>Acer x freemanii</i>	108	43	12	12	2	0	0	0	177	0.58
<i>Gymnocladus dioica</i>	132	6	4	10	18	5	1	0	176	0.57
<i>Morus alba</i>	5	3	15	47	44	28	24	6	172	0.56
<i>Fraxinus angustifolia</i> ssp.										
<i>Oxycarpa</i>	9	12	30	65	39	13	1	0	169	0.55
<i>Magnolia grandiflora</i>	20	61	31	44	4	0	0	0	160	0.52
<i>Pinus halepensis</i>	6	8	13	27	22	26	35	20	157	0.51
<i>Quercus buckleyi</i>	145	7	1	0	0	0	0	0	153	0.50
<i>Pistacia</i> spp.	151	0	0	1	0	0	0	0	152	0.50
<i>Quercus ilex</i>	22	20	16	28	41	23	1	1	152	0.50
<i>Ulmus</i> spp.	117	9	2	4	5	0	4	4	145	0.47
<i>Quercus canbyi</i>	143	0	0	0	0	0	0	0	143	0.47
<i>Melia azedarach</i>	6	8	16	31	31	18	14	1	125	0.41
<i>Fraxinus</i> spp.	14	32	27	31	14	5	1	0	124	0.40
<i>Prunus</i> spp.	9	50	26	23	7	5	1	1	122	0.40
<i>x Chitalpa tashkentensis</i>	73	28	9	2	1	0	0	0	113	0.37
<i>Tilia cordata</i>	12	18	36	23	4	2	0	0	95	0.31
<i>Sophora japonica</i>	4	0	1	10	49	25	1	0	90	0.29
<i>Quercus shumardii</i>	66	6	10	3	0	0	1	0	86	0.28
<i>Populus fremontii</i>	4	14	10	24	15	10	5	3	85	0.28
<i>Acer platanooides</i>	8	64	11	1	0	0	0	0	84	0.27
<i>Pinus</i> spp.	0	2	3	41	22	12	2	0	82	0.27
<i>Robinia pseudoacacia</i>	4	21	23	26	4	2	2	0	82	0.27
<i>Quercus coccinea</i>	42	11	14	10	1	2	0	0	80	0.26
<i>Ligustrum lucidum</i>	3	15	10	41	6	1	0	2	78	0.25
<i>Chilopsis linearis</i>	69	4	2	1	0	1	0	0	77	0.25
<i>Quercus robur</i>	62	4	2	7	2	0	0	0	77	0.25
<i>Prunus serrulata</i>	18	23	23	8	0	1	0	0	73	0.24
<i>Searsia lancea</i>	1	0	5	11	28	19	1	1	66	0.22
<i>Carpinus betulus</i>	16	2	12	27	7	0	0	0	64	0.21
<i>Juglans regia</i>	2	3	3	39	7	5	3	1	63	0.21
<i>Cupressus arizonica</i>	8	9	12	23	7	0	0	0	59	0.19
<i>Maclura pomifera</i>	56	0	0	0	0	0	0	0	56	0.18
<i>Pinus pinea</i>	0	1	1	5	6	28	10	4	55	0.18
<i>Ostrya virginiana</i>	2	27	17	4	1	1	0	0	52	0.17
<i>Populus nigra</i> v. <i>italica</i>	0	9	12	27	3	0	0	1	52	0.17
<i>Chionanthus retusus</i>	43	0	1	0	0	0	0	0	44	0.14
<i>Schinus molle</i>	0	3	8	14	11	8	0	0	44	0.14
<i>Acer campestre</i>	0	3	9	29	1	1	0	0	43	0.14
<i>Acacia melanoxylon</i>	4	0	5	16	9	6	1	0	41	0.13
<i>Salix</i> spp.	0	0	8	18	9	3	0	2	40	0.13
<i>Pinus nigra</i>	0	7	26	5	0	0	0	0	38	0.12
<i>Eucalyptus camaldulensis</i>	2	1	2	12	2	8	5	4	36	0.12

Species	DBH Class (inches)								# of Trees	% of Pop.
	0 - 4	4 - 8	8 - 12	12 - 18	18 - 24	24 - 30	30 - 36	>36		
<i>Liquidambar styraciflua</i>	0	1	4	22	5	1	0	0	33	0.11
<i>Acer saccharinum</i>	1	4	3	7	7	6	2	1	31	0.10
<i>Aesculus californica</i>	10	14	2	4	1	0	0	0	31	0.10
<i>Calocedrus decurrens</i>	4	7	2	4	4	7	1	2	31	0.10
<i>Albizia julibrissin</i>	4	1	1	18	6	0	0	0	30	0.10
<i>Chamaecyparis</i>	1	7	5	8	5	3	0	0	29	0.09
<i>Quercus macrocarpa</i>	8	6	2	2	5	4	2	0	29	0.09
<i>Geijera parviflora</i>	0	8	8	6	1	2	1	1	27	0.09
<i>Robinia spp.</i>	1	4	11	7	2	0	0	1	26	0.08
<i>Umbellularia californica</i>	0	3	2	12	8	0	1	0	26	0.08
<i>Chilopsis spp.</i>	25	0	0	0	0	0	0	0	25	0.08
<i>Platanus occidentalis</i>	3	1	4	5	6	3	2	0	24	0.08
<i>Fraxinus ornus</i>	0	2	3	10	7	1	0	0	23	0.07
<i>Prunus dulcis</i>	7	6	4	3	2	1	0	0	23	0.07
<i>Quercus spp.</i>	8	3	1	9	1	1	0	0	23	0.07
<i>Tilia tomentosa</i>	13	8	1	0	0	0	0	0	22	0.07
<i>Betula pendula</i>	4	6	8	2	1	0	0	0	21	0.07
<i>Quercus frainetto</i>	17	2	2	0	0	0	0	0	21	0.07
<i>Pinus thunbergii</i>	0	1	8	9	2	0	0	0	20	0.07
<i>Casuarina stricta</i>	0	0	0	0	4	7	4	4	19	0.06
<i>Parrotia persica</i>	19	0	0	0	0	0	0	0	19	0.06
<i>Tilia americana</i>	3	2	5	4	3	0	1	0	18	0.06
<i>Ulmus americana</i>	5	0	0	0	1	2	5	5	18	0.06
<i>Eucalyptus polyanthemos</i>	0	1	3	3	2	4	3	1	17	0.06
<i>Ficus spp.</i>	4	2	3	5	2	0	1	0	17	0.06
<i>Platanus orientalis</i>	3	0	3	9	2	0	0	0	17	0.06
<i>Cinnamomum camphora</i>	0	4	7	1	2	1	1	0	16	0.05
<i>Taxodium distichum</i>	16	0	0	0	0	0	0	0	16	0.05
<i>Catalpa speciosa</i>	0	2	6	5	1	1	0	0	15	0.05
<i>Prosopis chilensis</i>	15	0	0	0	0	0	0	0	15	0.05
<i>Prunus persica</i>	3	3	1	7	1	0	0	0	15	0.05
<i>Tilia x euchlora</i>	0	0	4	11	0	0	0	0	15	0.05
<i>Acer palmatum</i>	10	4	0	0	0	0	0	0	14	0.05
<i>Ceratonia siliqua</i>	1	1	1	1	2	6	1	1	14	0.05
<i>Juglans nigra</i>	1	1	0	3	2	0	1	6	14	0.05
<i>Maytenus boaria</i>	2	0	1	4	3	1	1	1	13	0.04
<i>Pinus sylvestris</i>	1	1	2	4	2	3	0	0	13	0.04
<i>Paulownia tomentosa</i>	2	1	2	3	4	0	0	0	12	0.04
<i>Tristaniopsis spp.</i>	9	2	1	0	0	0	0	0	12	0.04
<i>Acer negundo</i>	3	4	2	0	2	0	0	0	11	0.04
<i>Afrocarpus gracilior</i>	0	2	8	1	0	0	0	0	11	0.04
<i>Alnus cordata</i>	0	0	4	6	1	0	0	0	11	0.04
<i>Cedrus atlantica</i>	1	1	3	3	2	1	0	0	11	0.04
<i>Quercus cerris</i>	1	0	0	2	3	2	3	0	11	0.04

Species	DBH Class (inches)								# of Trees	% of Pop.
	0 - 4	4 - 8	8 - 12	12 - 18	18 - 24	24 - 30	30 - 36	>36		
<i>Alnus rhombifolia</i>	1	3	0	4	2	0	0	0	10	0.03
<i>Eriobotrya japonica</i>	3	3	3	1	0	0	0	0	10	0.03
<i>Quercus palustris</i>	5	4	1	0	0	0	0	0	10	0.03
<i>Acer truncatum x platanoides</i>	2	4	0	3	0	0	0	0	9	0.03
<i>Eucalyptus melliodora</i>	0	0	1	1	0	5	2	0	9	0.03
<i>Prunus Kanzan</i>	6	1	1	1	0	0	0	0	9	0.03
<i>Pterocarya stenoptera</i>	0	2	0	1	4	0	0	2	9	0.03
<i>Punica spp.</i>	1	1	3	2	2	0	0	0	9	0.03
<i>Quercus laurifolia</i>	0	1	4	4	0	0	0	0	9	0.03
<i>Ulmus procera</i>	0	0	0	1	2	0	2	4	9	0.03
<i>Magnolia x soulangeana</i>	4	0	3	1	0	0	0	0	8	0.03
<i>Melaleuca linariifolia</i>	0	1	0	2	4	1	0	0	8	0.03
<i>Ailanthus altissima</i>	1	1	1	1	2	0	1	0	7	0.02
<i>Crataegus spp.</i>	2	3	0	2	0	0	0	0	7	0.02
<i>Eucalyptus spp.</i>	0	0	0	1	1	2	0	3	7	0.02
<i>Quercus chrysolepis</i>	7	0	0	0	0	0	0	0	7	0.02
<i>Acer pseudoplatanus</i>	1	0	1	4	0	0	0	0	6	0.02
<i>Eucalyptus nicholii</i>	0	0	1	2	0	1	2	0	6	0.02
<i>Quercus muehlenbergii</i>	5	0	0	1	0	0	0	0	6	0.02
<i>Salix babylonica</i>	0	0	0	0	0	1	5	0	6	0.02
<i>Betula nigra</i>	3	1	1	0	0	0	0	0	5	0.02
<i>Carpinus caroliniana</i>	2	3	0	0	0	0	0	0	5	0.02
<i>Quercus imbricaria</i>	5	0	0	0	0	0	0	0	5	0.02
<i>Xylosma spp.</i>	0	4	1	0	0	0	0	0	5	0.02
<i>Cupressus sempervirens</i>	0	0	2	2	0	0	0	0	4	0.01
<i>Lagerstroemia subcostata</i>	4	0	0	0	0	0	0	0	4	0.01
<i>Magnoliopsida</i>	4	0	0	0	0	0	0	0	4	0.01
<i>Myrtuss pp.</i>	0	0	2	0	0	2	0	0	4	0.01
<i>Picea pungens</i>	1	1	1	1	0	0	0	0	4	0.01
<i>Thuja plicata</i>	0	0	0	3	0	1	0	0	4	0.01
<i>Ulmus pumila</i>	0	0	0	0	0	0	1	3	4	0.01
<i>Acer griseum</i>	2	1	0	0	0	0	0	0	3	0.01
<i>Brachychiton populneus</i>	0	0	1	1	1	0	0	0	3	0.01
<i>Eucalyptus globulus</i>	0	0	0	1	0	1	1	0	3	0.01
<i>Juglans spp.</i>	0	1	0	1	0	0	0	1	3	0.01
<i>Metasequoia glyptostroboides</i>	3	0	0	0	0	0	0	0	3	0.01
<i>Parkinsonia aculeata</i>	3	0	0	0	0	0	0	0	3	0.01
<i>Pinus radiata</i>	0	0	1	0	1	1	0	0	3	0.01
<i>Acer spp.</i>	0	0	0	2	0	0	0	0	2	0.01
<i>Acer saccharum</i>	0	0	1	1	0	0	0	0	2	0.01
<i>Callistemon spp.</i>	2	0	0	0	0	0	0	0	2	0.01
<i>Citrus spp.</i>	0	0	0	1	1	0	0	0	2	0.01
<i>Citrus limon</i>	0	2	0	0	0	0	0	0	2	0.01

Species	DBH Class (inches)								# of Trees	% of Pop.
	0 - 4	4 - 8	8 - 12	12 - 18	18 - 24	24 - 30	30 - 36	>36		
<i>Corymbia maculata</i>	0	1	0	0	0	0	1	0	2	0.01
<i>Crataegus phaenopyrum</i>	2	0	0	0	0	0	0	0	2	0.01
<i>Koelreuteria elegans</i>	1	1	0	0	0	0	0	0	2	0.01
<i>Photinia spp.</i>	0	1	1	0	0	0	0	0	2	0.01
<i>Pinus coulteri</i>	0	0	0	2	0	0	0	0	2	0.01
<i>Prunus armeniaca</i>	0	1	1	0	0	0	0	0	2	0.01
<i>Prunus lyonii</i>	1	0	1	0	0	0	0	0	2	0.01
<i>Prunus x yedoensis</i>	2	0	0	0	0	0	0	0	2	0.01
<i>Quercus castaneifolia</i>	2	0	0	0	0	0	0	0	2	0.01
<i>Quercus nigra</i>	1	0	0	1	0	0	0	0	2	0.01
<i>Quercus phellos</i>	1	0	0	0	0	0	0	1	2	0.01
<i>Washingtonia filifera</i>	2	0	0	0	0	0	0	0	2	0.01
<i>Acer macrophyllum</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Brachychiton acerifolius</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Carya illinoensis</i>	0	0	1	0	0	0	0	0	1	0.00
<i>Cercidiphyllum japonicum</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Elaeocarpus decipiens</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Firmiana simplex</i>	0	0	0	1	0	0	0	0	1	0.00
<i>Juniperus virginiana</i>	0	0	1	0	0	0	0	0	1	0.00
<i>Larix occidentalis</i>	0	1	0	0	0	0	0	0	1	0.00
<i>Magnolia virginiana</i>	0	0	1	0	0	0	0	0	1	0.00
<i>Malus x arnoldiana</i>	0	0	1	0	0	0	0	0	1	0.00
<i>Nyssa sylvatica</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Ostrya carpinifolia</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Persea spp.</i>	1	0	0	0	0	0	0	0	1	0.00
<i>Pistacia vera</i>	0	0	0	1	0	0	0	0	1	0.00
<i>Prunus subhirtella</i>	0	0	0	0	0	1	0	0	1	0.00
<i>Prunus x blireiana</i>	0	0	0	1	0	0	0	0	1	0.00
<i>Pyrus betulifolia</i>	0	1	0	0	0	0	0	0	1	0.00
<i>Quercus engelmannii</i>	0	0	0	1	0	0	0	0	1	0.00
<i>Tetradium spp.</i>	0	0	0	0	1	0	0	0	1	0.00
<i>Thuja spp.</i>	0	0	0	0	0	0	0	1	1	0.00
<i>Ulmus x hollandica</i>	0	0	0	0	0	0	0	1	1	0.00
<i>Washingtonia spp.</i>	0	0	0	1	0	0	0	0	1	0.00



**Table 16: Importance Value (IV) for All Tree Species**

Species	% of Pop.	% of Leaf Area	Importance Value (IV)
<i>Platanus x acerifolia</i>	7.75	11.30	19.10
<i>Pistacia chinensis</i>	6.57	3.50	10.10
<i>Lagerstroemia indica</i>	6.25	0.80	7.00
<i>Quercus lobata</i>	5.97	7.00	13.00
<i>Sequoia sempervirens</i>	4.95	6.60	11.60
<i>Pyrus calleryana</i>	4.84	3.40	8.20
<i>Triadica sebifera</i>	4.31	3.80	8.10
<i>Celtis sinensis</i>	3.35	6.10	9.40
<i>Fraxinus holotricha</i>	2.67	5.00	7.70
<i>Vitex agnus-castus</i>	2.56	2.40	5.00
<i>Casuarina cunninghamiana</i>	2.15	1.70	3.90
<i>Pinus canariensis</i>	1.84	4.00	5.90
<i>Celtis australis</i>	1.76	3.00	4.80
<i>Ulmus parvifolia</i>	1.71	0.90	2.60
<i>Ginkgo biloba</i>	1.65	0.60	2.30
<i>Acer buergerianum</i>	1.40	0.70	2.10
<i>Koelreuteria paniculata</i>	1.34	1.60	3.00
<i>Pinus brutia</i>	1.23	1.70	2.90
<i>Gleditsia triacanthos</i>	1.18	1.40	2.60
<i>Quercus douglasii</i>	1.18	0.20	1.40
<i>Fraxinus velutina</i>	1.15	1.90	3.10
<i>Quercus suber</i>	1.06	2.00	3.00
<i>Fraxinus americana</i>	1.04	0.50	1.60
<i>Platanus racemosa</i>	1.02	1.90	2.90
<i>Koelreuteria bipinnata</i>	0.99	0.30	1.30
<i>Quercus virginiana</i>	0.91	1.10	2.00
<i>Cedrus deodara</i>	0.89	0.90	1.80
<i>Acer rubrum</i>	0.84	0.10	1.00
<i>Cercis canadensis</i>	0.79	0.30	1.10
<i>Quercus agrifolia</i>	0.77	1.20	1.90
<i>Ulmus davidiana v. japonica</i>	0.76	0.10	0.90
<i>Olea europaea</i>	0.74	1.00	1.70
<i>Celtis occidentalis</i>	0.72	0.90	1.60
<i>Fraxinus uhdei</i>	0.70	1.10	1.80
<i>Zelkova serrata</i>	0.69	0.20	0.90
<i>Juglans hindsii</i>	0.65	1.00	1.70
<i>Laurus nobilis</i>	0.65	0.40	1.10
<i>Malus spp.</i>	0.63	0.10	0.70
<i>Quercus wislizeni</i>	0.62	0.80	1.50
<i>Prunus cerasifera</i>	0.61	0.30	0.90
<i>Arbutus unedo</i>	0.61	0.30	0.90
<i>Liriodendron tulipifera</i>	0.61	0.80	1.40

Species	% of Pop.	% of Leaf Area	Importance Value (IV)
<i>Quercus rubra</i>	0.60	0.10	0.70
<i>Eucalyptus sideroxylon</i>	0.58	1.50	2.10
<i>Acer x freemanii</i>	0.58	0.30	0.80
<i>Gymnocladus dioica</i>	0.57	0.20	0.80
<i>Morus alba</i>	0.56	0.80	1.40
<i>Fraxinus angustifolia ssp. Oxycarpa</i>	0.55	0.90	1.40
<i>Magnolia grandiflora</i>	0.52	0.30	0.90
<i>Pinus halepensis</i>	0.51	1.00	1.60
<i>Quercus buckleyi</i>	0.50	0.00	0.50
<i>Pistacia spp.</i>	0.50	0.00	0.50
<i>Quercus ilex</i>	0.50	1.20	1.70
<i>Ulmus spp.</i>	0.47	0.10	0.60
<i>Quercus canbyi</i>	0.47	0.00	0.50
<i>Melia azedarach</i>	0.41	0.60	1.00
<i>Fraxinus spp.</i>	0.40	0.40	0.80
<i>Prunus spp.</i>	0.40	0.20	0.60
<i>x Chitalpa tashkentensis</i>	0.37	0.00	0.40
<i>Tilia cordata</i>	0.31	0.30	0.60
<i>Sophora japonica</i>	0.29	0.40	0.70
<i>Quercus shumardii</i>	0.28	0.10	0.40
<i>Populus fremontii</i>	0.28	0.40	0.60
<i>Acer platanoides</i>	0.27	0.10	0.30
<i>Pinus spp.</i>	0.27	0.30	0.60
<i>Robinia pseudoacacia</i>	0.27	0.20	0.40
<i>Quercus coccinea</i>	0.26	0.20	0.40
<i>Ligustrum lucidum</i>	0.25	0.30	0.60
<i>Chilopsis linearis</i>	0.25	0.00	0.30
<i>Quercus robur</i>	0.25	0.10	0.30
<i>Prunus serrulata</i>	0.24	0.10	0.30
<i>Searsia lancea</i>	0.22	0.20	0.40
<i>Carpinus betulus</i>	0.21	0.30	0.50
<i>Juglans regia</i>	0.21	0.30	0.50
<i>Cupressus arizonica</i>	0.19	0.10	0.30
<i>Maclura pomifera</i>	0.18	0.00	0.20
<i>Pinus pinea</i>	0.18	0.40	0.60
<i>Ostrya virginiana</i>	0.17	0.10	0.30
<i>Populus nigra v. italica</i>	0.17	0.20	0.30
<i>Chionanthus retusus</i>	0.14	0.00	0.10
<i>Schinus molle</i>	0.14	0.10	0.30
<i>Acer campestre</i>	0.14	0.20	0.40
<i>Acacia melanoxylon</i>	0.13	0.30	0.40
<i>Salix spp.</i>	0.13	0.10	0.30
<i>Pinus nigra</i>	0.12	0.00	0.20

Species	% of Pop.	% of Leaf Area	Importance Value (IV)
<i>Eucalyptus camaldulensis</i>	0.12	0.60	0.70
<i>Liquidambar styraciflua</i>	0.11	0.20	0.30
<i>Acer saccharinum</i>	0.10	0.20	0.30
<i>Aesculus californica</i>	0.10	0.10	0.20
<i>Calocedrus decurrens</i>	0.10	0.10	0.20
<i>Albizia julibrissin</i>	0.10	0.10	0.20
<i>Chamaecyparis</i>	0.09	0.10	0.20
<i>Quercus macrocarpa</i>	0.09	0.20	0.30
<i>Geijera parviflora</i>	0.09	0.00	0.10
<i>Robinia spp.</i>	0.08	0.10	0.20
<i>Umbellularia californica</i>	0.08	0.10	0.20
<i>Chilopsis spp.</i>	0.08	0.00	0.10
<i>Platanus occidentalis</i>	0.08	0.20	0.30
<i>Fraxinus ornus</i>	0.07	0.10	0.20
<i>Prunus dulcis</i>	0.07	0.00	0.10
<i>Quercus spp.</i>	0.07	0.10	0.10
<i>Tilia tomentosa</i>	0.07	0.00	0.10
<i>Betula pendula</i>	0.07	0.00	0.10
<i>Quercus frainetto</i>	0.07	0.00	0.10
<i>Pinus thunbergii</i>	0.07	0.10	0.10
<i>Casuarina stricta</i>	0.06	0.10	0.10
<i>Parrotia persica</i>	0.06	0.00	0.10
<i>Tilia americana</i>	0.06	0.10	0.10
<i>Ulmus americana</i>	0.06	0.10	0.20
<i>Eucalyptus polyanthemos</i>	0.06	0.20	0.30
<i>Ficus spp.</i>	0.06	0.00	0.10
<i>Platanus orientalis</i>	0.06	0.10	0.10
<i>Cinnamomum camphora</i>	0.05	0.00	0.10
<i>Taxodium distichum</i>	0.05	0.00	0.10
<i>Catalpa speciosa</i>	0.05	0.00	0.10
<i>Prosopis chilensis</i>	0.05	0.00	0.00
<i>Prunus persica</i>	0.05	0.00	0.10
<i>Tilia x euchlora</i>	0.05	0.10	0.10
<i>Acer palmatum</i>	0.05	0.00	0.00
<i>Ceratonia siliqua</i>	0.05	0.10	0.10
<i>Juglans nigra</i>	0.05	0.10	0.10
<i>Maytenus boaria</i>	0.04	0.00	0.10
<i>Pinus sylvestris</i>	0.04	0.10	0.10
<i>Paulownia tomentosa</i>	0.04	0.00	0.10
<i>Tristaniopsis spp.</i>	0.04	0.00	0.00
<i>Acer negundo</i>	0.04	0.00	0.10
<i>Afrocarpus gracilior</i>	0.04	0.00	0.10
<i>Alnus cordata</i>	0.04	0.10	0.10

Species	% of Pop.	% of Leaf Area	Importance Value (IV)
<i>Cedrus atlantica</i>	0.04	0.00	0.10
<i>Quercus cerris</i>	0.04	0.10	0.10
<i>Alnus rhombifolia</i>	0.03	0.00	0.10
<i>Eriobotrya japonica</i>	0.03	0.00	0.00
<i>Quercus palustris</i>	0.03	0.00	0.00
<i>Acer truncatum x platanoides</i>	0.03	0.00	0.10
<i>Eucalyptus melliodora</i>	0.03	0.10	0.20
<i>Prunus Kanzan</i>	0.03	0.00	0.00
<i>Pterocarya stenoptera</i>	0.03	0.00	0.10
<i>Punica spp.</i>	0.03	0.00	0.10
<i>Quercus laurifolia</i>	0.03	0.00	0.10
<i>Ulmus procera</i>	0.03	0.00	0.10
<i>Magnolia x soulangeana</i>	0.03	0.00	0.00
<i>Melaleuca linariifolia</i>	0.03	0.00	0.10
<i>Ailanthus altissima</i>	0.02	0.00	0.00
<i>Crataegus spp.</i>	0.02	0.00	0.00
<i>Eucalyptus spp.</i>	0.02	0.10	0.20
<i>Quercus chrysolepis</i>	0.02	0.00	0.00
<i>Acer pseudoplatanus</i>	0.02	0.00	0.00
<i>Eucalyptus nicholii</i>	0.02	0.10	0.10
<i>Quercus muehlenbergii</i>	0.02	0.00	0.00
<i>Salix babylonica</i>	0.02	0.00	0.10
<i>Betula nigra</i>	0.02	0.00	0.00
<i>Carpinus caroliniana</i>	0.02	0.00	0.00
<i>Quercus imbricaria</i>	0.02	0.00	0.00
<i>Xylosma spp.</i>	0.02	0.00	0.00
<i>Cupressus sempervirens</i>	0.01	0.00	0.00
<i>Lagerstroemia subcostata</i>	0.01	0.00	0.00
<i>Magnoliopsida</i>	0.01	0.00	0.00
<i>Myrtuss pp.</i>	0.01	0.00	0.10
<i>Picea pungens</i>	0.01	0.00	0.00
<i>Thuja plicata</i>	0.01	0.00	0.00
<i>Ulmus pumila</i>	0.01	0.00	0.00
<i>Acer griseum</i>	0.01	0.00	0.00
<i>Brachychiton populneus</i>	0.01	0.00	0.00
<i>Eucalyptus globulus</i>	0.01	0.00	0.10
<i>Juglans spp.</i>	0.01	0.00	0.00
<i>Metasequoia glyptostroboides</i>	0.01	0.00	0.00
<i>Parkinsonia aculeata</i>	0.01	0.00	0.00
<i>Pinus radiata</i>	0.01	0.00	0.00
<i>Acer spp.</i>	0.01	0.00	0.00
<i>Acer saccharum</i>	0.01	0.00	0.00
<i>Callistemon spp.</i>	0.01	0.00	0.00

Species	% of Pop.	% of Leaf Area	Importance Value (IV)
<i>Citrus spp.</i>	0.01	0.00	0.00
<i>Citrus limon</i>	0.01	0.00	0.00
<i>Corymbia maculata</i>	0.01	0.00	0.00
<i>Crataegus phaenopyrum</i>	0.01	0.00	0.00
<i>Koelreuteria elegans</i>	0.01	0.00	0.00
<i>Photinia spp.</i>	0.01	0.00	0.00
<i>Pinus coulteri</i>	0.01	0.00	0.00
<i>Prunus armeniaca</i>	0.01	0.00	0.00
<i>Prunus lyonii</i>	0.01	0.00	0.00
<i>Prunus x yedoensis</i>	0.01	0.00	0.00
<i>Quercus castaneifolia</i>	0.01	0.00	0.00
<i>Quercus nigra</i>	0.01	0.00	0.00
<i>Quercus phellos</i>	0.01	0.00	0.00
<i>Washingtonia filifera</i>	0.01	0.00	0.00
<i>Acer macrophyllum</i>	0.00	0.00	0.00
<i>Brachychiton acerifolius</i>	0.00	0.00	0.00
<i>Carya illinoensis</i>	0.00	0.00	0.00
<i>Cercidiphyllum japonicum</i>	0.00	0.00	0.00
<i>Elaeocarpus decipiens</i>	0.00	0.00	0.00
<i>Firmiana simplex</i>	0.00	0.00	0.00
<i>Juniperus virginiana</i>	0.00	0.00	0.00
<i>Larix occidentalis</i>	0.00	0.00	0.00
<i>Magnolia virginiana</i>	0.00	0.00	0.00
<i>Malus x arnoldiana</i>	0.00	0.00	0.00
<i>Nyssa sylvatica</i>	0.00	0.00	0.00
<i>Ostrya carpinifolia</i>	0.00	0.00	0.00
<i>Persea spp.</i>	0.00	0.00	0.00
<i>Pistacia vera</i>	0.00	0.00	0.00
<i>Prunus subhirtella</i>	0.00	0.00	0.00
<i>Prunus x blireiana</i>	0.00	0.00	0.00
<i>Pyrus betulifolia</i>	0.00	0.00	0.00
<i>Quercus engelmannii</i>	0.00	0.00	0.00
<i>Tetradium spp.</i>	0.00	0.00	0.00
<i>Thuja spp.</i>	0.00	0.00	0.00
<i>Ulmus x hollandica</i>	0.00	0.00	0.00
<i>Washingtonia spp.</i>	0.00	0.00	0.00



**Table 17: Condition and RPI for All Tree Species**

Species	Excellent (%)	Good (%)	Fair (%)	Poor (%)	Dead (%)	RPI	Number of Trees	% of Pop.
<i>Platanus x acerifolia</i>	0.00	42.60	52.20	4.70	0.40	<b>0.98</b>	2,380	7.75
<i>Pistacia chinensis</i>	0.00	79.30	18.50	1.80	0.30	<b>1.09</b>	2,016	6.57
<i>Lagerstroemia indica</i>	0.00	59.00	36.20	4.60	0.20	<b>1.03</b>	1,918	6.25
<i>Quercus lobata</i>	0.00	46.00	49.40	3.90	0.60	<b>0.99</b>	1,833	5.97
<i>Sequoia sempervirens</i>	0.00	54.20	42.00	2.90	1.00	<b>1.01</b>	1,518	4.95
<i>Pyrus calleryana</i>	0.00	8.40	65.60	24.10	2.00	<b>0.81</b>	1,484	4.84
<i>Triadica sebifera</i>	0.00	54.50	42.10	3.10	0.40	<b>1.02</b>	1,322	4.31
<i>Celtis sinensis</i>	0.20	55.40	38.10	5.50	0.80	<b>1.01</b>	1,028	3.35
<i>Fraxinus holotricha</i>	0.00	31.40	60.10	8.10	0.40	<b>0.94</b>	818	2.67
<i>Vitex agnus-castus</i>	0.00	77.50	19.90	2.50	0.00	<b>1.09</b>	787	2.56
<i>Casuarina cunninghamiana</i>	0.00	71.60	26.80	1.20	0.50	<b>1.07</b>	661	2.15
<i>Pinus canariensis</i>	0.00	73.60	22.30	3.90	0.20	<b>1.07</b>	564	1.84
<i>Celtis australis</i>	0.00	83.20	14.80	1.50	0.60	<b>1.10</b>	541	1.76
<i>Ulmus parvifolia</i>	0.00	78.70	18.90	1.90	0.60	<b>1.09</b>	525	1.71
<i>Ginkgo biloba</i>	0.20	88.60	7.30	1.60	2.40	<b>1.10</b>	507	1.65
<i>Acer buergerianum</i>	0.00	90.30	9.00	0.50	0.20	<b>1.13</b>	431	1.40
<i>Koelreuteria paniculata</i>	0.20	66.50	27.70	5.10	0.50	<b>1.05</b>	412	1.34
<i>Pinus brutia</i>	0.00	12.90	78.40	7.90	0.80	<b>0.88</b>	379	1.23
<i>Gleditsia triacanthos</i>	0.00	19.60	71.10	9.40	0.00	<b>0.90</b>	363	1.18
<i>Quercus douglasii</i>	0.00	77.90	21.00	0.60	0.60	<b>1.09</b>	362	1.18
<i>Fraxinus velutina</i>	0.00	5.60	66.70	27.10	0.60	<b>0.80</b>	354	1.15
<i>Quercus suber</i>	0.00	68.90	26.20	3.70	1.20	<b>1.05</b>	325	1.06
<i>Fraxinus americana</i>	0.00	60.00	31.60	7.50	0.90	<b>1.02</b>	320	1.04
<i>Platanus racemosa</i>	0.00	27.20	66.10	5.80	1.00	<b>0.92</b>	313	1.02
<i>Koelreuteria bipinnata</i>	0.00	79.00	18.70	2.00	0.30	<b>1.09</b>	305	0.99
<i>Quercus virginiana</i>	0.00	75.50	20.50	4.00	0.00	<b>1.08</b>	278	0.91
<i>Cedrus deodara</i>	0.70	63.10	32.80	2.60	0.70	<b>1.04</b>	274	0.89
<i>Acer rubrum</i>	0.00	88.80	7.80	1.90	1.60	<b>1.11</b>	258	0.84
<i>Cercis canadensis</i>	0.00	51.60	42.60	4.50	1.20	<b>1.00</b>	244	0.79
<i>Quercus agrifolia</i>	0.00	59.30	37.30	1.70	1.70	<b>1.02</b>	236	0.77
<i>Ulmus davidiana v. japonica</i>	0.00	87.90	11.60	0.40	0.00	<b>1.12</b>	232	0.76
<i>Olea europaea</i>	0.00	15.50	81.90	2.70	0.00	<b>0.91</b>	226	0.74
<i>Celtis occidentalis</i>	0.00	27.00	66.70	5.90	0.50	<b>0.93</b>	222	0.72
<i>Fraxinus uhdei</i>	0.00	34.10	55.10	10.30	0.50	<b>0.94</b>	214	0.70
<i>Zelkova serrata</i>	0.00	86.80	7.50	5.70	0.00	<b>1.11</b>	212	0.69
<i>Juglans hindsii</i>	0.00	9.50	52.70	27.40	10.40	<b>0.73</b>	201	0.65
<i>Laurus nobilis</i>	0.00	24.20	70.70	4.50	0.50	<b>0.92</b>	198	0.65
<i>Malus spp.</i>	0.00	32.80	57.30	5.20	4.70	<b>0.91</b>	192	0.63
<i>Quercus wislizeni</i>	0.00	38.10	54.00	7.90	0.00	<b>0.96</b>	189	0.62
<i>Prunus cerasifera</i>	0.00	19.70	75.50	2.10	2.70	<b>0.90</b>	188	0.61
<i>Arbutus unedo</i>	0.00	40.10	54.00	3.20	2.70	<b>0.95</b>	187	0.61

Species	Excellent (%)	Good (%)	Fair (%)	Poor (%)	Dead (%)	RPI	Number of Trees	% of Pop.
<i>Liriodendron tulipifera</i>	0.00	51.90	35.80	9.60	2.70	<b>0.97</b>	187	0.61
<i>Quercus rubra</i>	0.00	79.20	15.30	1.60	3.80	<b>1.06</b>	183	0.60
<i>Eucalyptus sideroxylon</i>	0.00	5.10	91.60	3.40	0.00	<b>0.88</b>	178	0.58
<i>Acer x freemanii</i>	0.60	80.80	15.30	2.80	0.60	<b>1.09</b>	177	0.58
<i>Gymnocladus dioica</i>	0.00	88.10	10.20	1.10	0.60	<b>1.12</b>	176	0.57
<i>Morus alba</i>	0.00	20.90	69.80	9.30	0.00	<b>0.90</b>	172	0.56
<i>Fraxinus angustifolia ssp. Oxycarpa</i>	0.00	32.50	55.00	12.40	0.00	<b>0.93</b>	169	0.55
<i>Magnolia grandiflora</i>	0.00	23.10	61.90	15.00	0.00	<b>0.89</b>	160	0.52
<i>Pinus halepensis</i>	0.00	48.40	45.20	5.70	0.60	<b>0.99</b>	157	0.51
<i>Quercus buckleyi</i>	0.00	98.70	1.30	0.00	0.00	<b>1.16</b>	153	0.50
<i>Pistacia spp.</i>	0.00	99.30	0.70	0.00	0.00	<b>1.16</b>	152	0.50
<i>Quercus ilex</i>	0.00	71.70	25.00	1.30	2.00	<b>1.06</b>	152	0.50
<i>Ulmus spp.</i>	0.00	84.80	11.00	4.10	0.00	<b>1.11</b>	145	0.47
<i>Quercus canbyi</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	143	0.47
<i>Melia azedarach</i>	0.00	74.40	24.80	0.80	0.00	<b>1.08</b>	125	0.41
<i>Fraxinus spp.</i>	0.00	19.40	79.00	1.60	0.00	<b>0.92</b>	124	0.40
<i>Prunus spp.</i>	0.00	4.10	63.90	25.40	6.60	<b>0.75</b>	122	0.40
<i>x Chitalpa tashkentensis</i>	0.00	52.20	38.10	6.20	3.50	<b>0.97</b>	113	0.37
<i>Tilia cordata</i>	0.00	55.80	38.90	5.30	0.00	<b>1.02</b>	95	0.31
<i>Sophora japonica</i>	0.00	13.30	86.70	0.00	0.00	<b>0.91</b>	90	0.29
<i>Quercus shumardii</i>	0.00	68.60	24.40	2.30	4.70	<b>1.02</b>	86	0.28
<i>Populus fremontii</i>	0.00	16.50	68.20	7.10	8.20	<b>0.83</b>	85	0.28
<i>Acer platanoides</i>	0.00	27.40	51.20	14.30	7.10	<b>0.85</b>	84	0.27
<i>Pinus spp.</i>	0.00	23.20	50.00	26.80	0.00	<b>0.86</b>	82	0.27
<i>Robinia pseudoacacia</i>	0.00	14.60	75.60	6.10	3.70	<b>0.86</b>	82	0.27
<i>Quercus coccinea</i>	0.00	67.50	25.00	3.80	3.80	<b>1.02</b>	80	0.26
<i>Ligustrum lucidum</i>	0.00	12.80	84.60	2.60	0.00	<b>0.90</b>	78	0.25
<i>Chilopsis linearis</i>	0.00	93.50	6.50	0.00	0.00	<b>1.14</b>	77	0.25
<i>Quercus robur</i>	0.00	79.20	18.20	1.30	1.30	<b>1.09</b>	77	0.25
<i>Prunus serrulata</i>	0.00	50.70	39.70	1.40	8.20	<b>0.94</b>	73	0.24
<i>Searsia lancea</i>	0.00	12.10	84.80	0.00	3.00	<b>0.88</b>	66	0.22
<i>Carpinus betulus</i>	0.00	76.60	17.20	4.70	1.60	<b>1.07</b>	64	0.21
<i>Juglans regia</i>	0.00	4.80	69.80	17.50	7.90	<b>0.76</b>	63	0.21
<i>Cupressus arizonica</i>	0.00	37.30	37.30	22.00	3.40	<b>0.89</b>	59	0.19
<i>Maclura pomifera</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	56	0.18
<i>Pinus pinea</i>	0.00	54.50	45.50	0.00	0.00	<b>1.03</b>	55	0.18
<i>Ostrya virginiana</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	52	0.17
<i>Populus nigra v. italica</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	52	0.17
<i>Chionanthus retusus</i>	0.00	81.80	18.20	0.00	0.00	<b>1.11</b>	44	0.14
<i>Schinus molle</i>	0.00	15.90	84.10	0.00	0.00	<b>0.92</b>	44	0.14
<i>Acer campestre</i>	0.00	60.50	37.20	2.30	0.00	<b>1.04</b>	43	0.14
<i>Acacia melanoxylon</i>	0.00	61.00	36.60	2.40	0.00	<b>1.04</b>	41	0.13

Species	Excellent (%)	Good (%)	Fair (%)	Poor (%)	Dead (%)	RPI	Number of Trees	% of Pop.
<i>Salix spp.</i>	0.00	2.50	50.00	47.50	0.00	<b>0.74</b>	40	0.13
<i>Pinus nigra</i>	0.00	7.90	84.20	7.90	0.00	<b>0.87</b>	38	0.12
<i>Eucalyptus camaldulensis</i>	0.00	22.20	72.20	5.60	0.00	<b>0.92</b>	36	0.12
<i>Liquidambar styraciflua</i>	0.00	30.30	63.60	3.00	3.00	<b>0.92</b>	33	0.11
<i>Acer saccharinum</i>	0.00	38.70	48.40	9.70	3.20	<b>0.93</b>	31	0.10
<i>Aesculus californica</i>	0.00	71.00	25.80	0.00	3.20	<b>1.05</b>	31	0.10
<i>Calocedrus decurrens</i>	0.00	41.90	41.90	16.10	0.00	<b>0.95</b>	31	0.10
<i>Albizia julibrissin</i>	0.00	30.00	50.00	20.00	0.00	<b>0.90</b>	30	0.10
<i>Chamaecyparis</i>	0.00	20.70	55.20	24.10	0.00	<b>0.86</b>	29	0.09
<i>Quercus macrocarpa</i>	0.00	89.70	10.30	0.00	0.00	<b>1.13</b>	29	0.09
<i>Geijera parviflora</i>	0.00	18.50	81.50	0.00	0.00	<b>0.92</b>	27	0.09
<i>Robinia spp.</i>	0.00	19.20	65.40	15.40	0.00	<b>0.88</b>	26	0.08
<i>Umbellularia californica</i>	0.00	34.60	50.00	15.40	0.00	<b>0.93</b>	26	0.08
<i>Chilopsis spp.</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	25	0.08
<i>Platanus occidentalis</i>	0.00	29.20	66.70	4.20	0.00	<b>0.94</b>	24	0.08
<i>Fraxinus ornus</i>	0.00	47.80	43.50	8.70	0.00	<b>0.98</b>	23	0.07
<i>Prunus dulcis</i>	0.00	8.70	73.90	13.00	4.30	<b>0.82</b>	23	0.07
<i>Quercus spp.</i>	0.00	47.80	43.50	0.00	8.70	<b>0.93</b>	23	0.07
<i>Tilia tomentosa</i>	0.00	81.80	13.60	4.50	0.00	<b>1.10</b>	22	0.07
<i>Betula pendula</i>	0.00	9.50	61.90	23.80	4.80	<b>0.79</b>	21	0.07
<i>Quercus frainetto</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	21	0.07
<i>Pinus thunbergii</i>	0.00	35.00	35.00	30.00	0.00	<b>0.89</b>	20	0.07
<i>Casuarina stricta</i>	0.00	68.40	31.60	0.00	0.00	<b>1.07</b>	19	0.06
<i>Parrotia persica</i>	0.00	78.90	21.10	0.00	0.00	<b>1.10</b>	19	0.06
<i>Tilia americana</i>	0.00	77.80	22.20	0.00	0.00	<b>1.10</b>	18	0.06
<i>Ulmus americana</i>	0.00	50.00	33.30	16.70	0.00	<b>0.97</b>	18	0.06
<i>Eucalyptus polyanthemos</i>	0.00	5.90	94.10	0.00	0.00	<b>0.89</b>	17	0.06
<i>Ficus spp.</i>	0.00	29.40	41.20	17.60	11.80	<b>0.80</b>	17	0.06
<i>Platanus orientalis</i>	0.00	52.90	47.10	0.00	0.00	<b>1.02</b>	17	0.06
<i>Cinnamomum camphora</i>	0.00	6.30	75.00	12.50	6.30	<b>0.80</b>	16	0.05
<i>Taxodium distichum</i>	0.00	56.30	37.50	6.30	0.00	<b>1.02</b>	16	0.05
<i>Catalpa speciosa</i>	0.00	13.30	80.00	6.70	0.00	<b>0.89</b>	15	0.05
<i>Prosopis chilensis</i>	0.00	80.00	20.00	0.00	0.00	<b>1.10</b>	15	0.05
<i>Prunus persica</i>	0.00	20.00	80.00	0.00	0.00	<b>0.93</b>	15	0.05
<i>Tilia x euclora</i>	0.00	80.00	20.00	0.00	0.00	<b>1.10</b>	15	0.05
<i>Acer palmatum</i>	0.00	78.60	7.10	7.10	7.10	<b>1.02</b>	14	0.05
<i>Ceratonia siliqua</i>	0.00	28.60	71.40	0.00	0.00	<b>0.95</b>	14	0.05
<i>Juglans nigra</i>	0.00	35.70	21.40	35.70	7.10	<b>0.81</b>	14	0.05
<i>Maytenus boaria</i>	0.00	30.80	38.50	30.80	0.00	<b>0.87</b>	13	0.04
<i>Pinus sylvestris</i>	0.00	53.80	7.70	7.70	30.80	<b>0.74</b>	13	0.04
<i>Paulownia tomentosa</i>	0.00	0.00	50.00	50.00	0.00	<b>0.73</b>	12	0.04
<i>Tristaniopsis spp.</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	12	0.04
<i>Acer negundo</i>	0.00	9.10	45.50	18.20	27.30	<b>0.61</b>	11	0.04

Species	Excellent (%)	Good (%)	Fair (%)	Poor (%)	Dead (%)	RPI	Number of Trees	% of Pop.
<i>Afrocarpus gracilior</i>	0.00	90.90	9.10	0.00	0.00	<b>1.13</b>	11	0.04
<i>Alnus cordata</i>	0.00	45.50	54.50	0.00	0.00	<b>1.00</b>	11	0.04
<i>Cedrus atlantica</i>	0.00	54.50	36.40	9.10	0.00	<b>1.00</b>	11	0.04
<i>Quercus cerris</i>	0.00	72.70	27.30	0.00	0.00	<b>1.08</b>	11	0.04
<i>Alnus rhombifolia</i>	0.00	10.00	50.00	20.00	20.00	<b>0.67</b>	10	0.03
<i>Eriobotrya japonica</i>	0.00	60.00	40.00	0.00	0.00	<b>1.04</b>	10	0.03
<i>Quercus palustris</i>	0.00	80.00	20.00	0.00	0.00	<b>1.10</b>	10	0.03
<i>Acer truncatum x platanooides</i>	0.00	44.40	55.60	0.00	0.00	<b>1.00</b>	9	0.03
<i>Eucalyptus melliodora</i>	0.00	33.30	33.30	11.10	22.20	<b>0.74</b>	9	0.03
<i>Prunus Kanzan</i>	0.00	66.70	33.30	0.00	0.00	<b>1.06</b>	9	0.03
<i>Pterocarya stenoptera</i>	0.00	22.20	66.70	11.10	0.00	<b>0.90</b>	9	0.03
<i>Punica spp.</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	9	0.03
<i>Quercus laurifolia</i>	0.00	77.80	22.20	0.00	0.00	<b>1.10</b>	9	0.03
<i>Ulmus procera</i>	0.00	11.10	66.70	22.20	0.00	<b>0.84</b>	9	0.03
<i>Magnolia x soulangeana</i>	0.00	62.50	25.00	12.50	0.00	<b>1.02</b>	8	0.03
<i>Melaleuca linariifolia</i>	0.00	12.50	12.50	75.00	0.00	<b>0.69</b>	8	0.03
<i>Ailanthus altissima</i>	0.00	14.30	71.40	14.30	0.00	<b>0.87</b>	7	0.02
<i>Crataegus spp.</i>	0.00	42.90	14.30	42.90	0.00	<b>0.87</b>	7	0.02
<i>Eucalyptus spp.</i>	0.00	14.30	57.10	14.30	14.30	<b>0.75</b>	7	0.02
<i>Quercus chrysolepis</i>	0.00	57.10	0.00	0.00	42.90	<b>0.66</b>	7	0.02
<i>Acer pseudoplatanus</i>	0.00	66.70	16.70	0.00	16.70	<b>0.92</b>	6	0.02
<i>Eucalyptus nicholii</i>	0.00	50.00	50.00	0.00	0.00	<b>1.02</b>	6	0.02
<i>Quercus muehlenbergii</i>	0.00	83.30	16.70	0.00	0.00	<b>1.11</b>	6	0.02
<i>Salix babylonica</i>	0.00	16.70	66.70	16.70	0.00	<b>0.87</b>	6	0.02
<i>Betula nigra</i>	0.00	40.00	20.00	20.00	20.00	<b>0.75</b>	5	0.02
<i>Carpinus caroliniana</i>	0.00	80.00	20.00	0.00	0.00	<b>1.10</b>	5	0.02
<i>Quercus imbricaria</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	5	0.02
<i>Xylosma spp.</i>	0.00	60.00	20.00	20.00	0.00	<b>0.99</b>	5	0.02
<i>Cupressus sempervirens</i>	0.00	25.00	75.00	0.00	0.00	<b>0.94</b>	4	0.01
<i>Lagerstroemia subcostata</i>	0.00	75.00	25.00	0.00	0.00	<b>1.09</b>	4	0.01
<i>Magnoliopsida</i>	0.00	75.00	0.00	0.00	25.00	<b>0.87</b>	4	0.01
<i>Myrtuss pp.</i>	0.00	25.00	75.00	0.00	0.00	<b>0.94</b>	4	0.01
<i>Picea pungens</i>	0.00	25.00	75.00	0.00	0.00	<b>0.94</b>	4	0.01
<i>Thuja plicata</i>	0.00	0.00	75.00	25.00	0.00	<b>0.8</b>	4	0.01
<i>Ulmus pumila</i>	0.00	0.00	50.00	50.00	0.00	<b>0.73</b>	4	0.01
<i>Acer griseum</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	3	0.01
<i>Brachychiton populneus</i>	0.00	66.70	33.30	0.00	0.00	<b>1.06</b>	3	0.01
<i>Eucalyptus globulus</i>	0.00	66.70	0.00	33.30	0.00	<b>0.97</b>	3	0.01
<i>Juglans spp.</i>	0.00	33.30	33.30	33.30	0.00	<b>0.87</b>	3	0.01
<i>Metasequoia glyptostroboides</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	3	0.01
<i>Parkinsonia aculeata</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	3	0.01
<i>Pinus radiata</i>	0.00	33.30	66.70	0.00	0.00	<b>0.97</b>	3	0.01

Species	Excellent (%)	Good (%)	Fair (%)	Poor (%)	Dead (%)	RPI	Number of Trees	% of Pop.
<i>Acer spp.</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	2	0.01
<i>Acer saccharum</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	2	0.01
<i>Callistemon spp.</i>	0.00	50.00	50.00	0.00	0.00	<b>1.02</b>	2	0.01
<i>Citrus spp.</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	2	0.01
<i>Citrus limon</i>	0.00	50.00	50.00	0.00	0.00	<b>1.02</b>	2	0.01
<i>Corymbia maculata</i>	0.00	0.00	50.00	50.00	0.00	<b>0.73</b>	2	0.01
<i>Crataegus phaenopyrum</i>	0.00	50.00	50.00	0.00	0.00	<b>1.02</b>	2	0.01
<i>Koelreuteria elegans</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	2	0.01
<i>Photinia spp.</i>	0.00	0.00	50.00	0.00	50.00	<b>0.44</b>	2	0.01
<i>Pinus coulteri</i>	0.00	0.00	50.00	50.00	0.00	<b>0.73</b>	2	0.01
<i>Prunus armeniaca</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	2	0.01
<i>Prunus lyonii</i>	0.00	50.00	50.00	0.00	0.00	<b>1.02</b>	2	0.01
<i>Prunus x yedoensis</i>	0.00	50.00	50.00	0.00	0.00	<b>1.02</b>	2	0.01
<i>Quercus castaneifolia</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	2	0.01
<i>Quercus nigra</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	2	0.01
<i>Quercus phellos</i>	0.00	0.00	50.00	50.00	0.00	<b>0.73</b>	2	0.01
<i>Washingtonia filifera</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	2	0.01
<i>Acer macrophyllum</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Brachychiton acerifolius</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Carya illinoensis</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Cercidiphyllum japonicum</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Elaeocarpus decipiens</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Firmiana simplex</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Juniperus virginiana</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Larix occidentalis</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Magnolia virginiana</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Malus x arnoldiana</i>	0.00	0.00	0.00	0.00	100.00	<b>0</b>	1	0.00
<i>Nyssa sylvatica</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Ostrya carpinifolia</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Persea spp.</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Pistacia vera</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Prunus subhirtella</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Prunus x blireiana</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Pyrus betulifolia</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Quercus engelmannii</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00
<i>Tetradium spp.</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Thuja spp.</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Ulmus x hollandica</i>	0.00	0.00	100.00	0.00	0.00	<b>0.87</b>	1	0.00
<i>Washingtonia spp.</i>	0.00	100.00	0.00	0.00	0.00	<b>1.16</b>	1	0.00



**Table 18: Annual Benefits for All Species**

Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Avoided Runoff (gallon/yr.)	Avoided Runoff (\$/yr.)	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Platanus x acerifolia</i>	2,380	7.75	51.27	8,744.03	311,842.98	2,786.63	1.18	13,327.21
<i>Pistacia chinensis</i>	2,016	6.57	16.54	2,820.84	96,873.33	865.66	0.37	4,140.07
<i>Lagerstroemia indica</i>	1,918	6.25	9.87	1,682.83	21,258.37	189.96	0.08	908.52
<i>Quercus lobata</i>	1,833	5.97	28.02	4,778.93	192,640.22	1,721.43	0.73	8,232.85
<i>Sequoia sempervirens</i>	1,518	4.95	21.13	3,603.16	182,536.66	1,631.15	0.69	7,801.06
<i>Pyrus calleryana</i>	1,484	4.84	24.73	4,217.58	93,563.73	836.08	0.36	3,998.63
<i>Triadica sebifera</i>	1,322	4.31	40.65	6,932.12	105,529.27	943.01	0.40	4,510.00
<i>Celtis sinensis</i>	1,028	3.35	3.28	559.13	166,982.41	1,492.15	0.63	7,136.32
<i>Fraxinus holotricha</i>	818	2.67	18.35	3,129.21	137,307.49	1,226.98	0.52	5,868.10
<i>Vitex agnus-castus</i>	787	2.56	4.52	771.33	66,256.99	592.07	0.25	2,831.62
<i>Casuarina cunninghamiana</i>	661	2.15	30.51	5,203.75	46,852.52	418.67	0.18	2,002.33
<i>Pinus canariensis</i>	564	1.84	9.80	1,671.14	110,484.59	987.29	0.42	4,721.77
<i>Celtis australis</i>	541	1.76	1.63	277.64	82,603.12	738.14	0.31	3,530.20
<i>Ulmus parvifolia</i>	525	1.71	7.61	1,298.44	25,309.14	226.16	0.10	1,081.64
<i>Ginkgo biloba</i>	507	1.65	0.89	152.03	17,637.42	157.61	0.07	753.77
<i>Acer buergerianum</i>	431	1.40	2.72	464.13	19,967.53	178.43	0.08	853.35
<i>Koelreuteria paniculata</i>	412	1.34	4.95	844.62	44,940.65	401.59	0.17	1,920.63
<i>Pinus spp.</i>	379	1.23	6.17	1,051.66	46,404.81	414.67	0.18	1,983.20
<i>Gleditsia triacanthos</i>	363	1.18	8.36	1,426.45	38,275.43	342.03	0.15	1,635.78
<i>Quercus douglasii</i>	362	1.18	0.65	110.83	5,825.45	52.06	0.02	248.96
<i>Fraxinus velutina</i>	354	1.15	5.42	923.86	52,726.15	471.16	0.20	2,253.35
<i>Quercus suber</i>	325	1.06	4.43	755.23	54,695.39	488.76	0.21	2,337.51
<i>Fraxinus americana</i>	320	1.04	2.26	385.45	13,964.41	124.79	0.05	596.80
<i>Platanus racemosa</i>	313	1.02	2.62	447.07	50,972.86	455.49	0.19	2,178.43
<i>Koelreuteria bipinnata</i>	305	0.99	0.73	124.46	8,335.02	74.48	0.03	356.21
<i>Quercus virginiana</i>	278	0.91	7.45	1,270.65	28,912.73	258.36	0.11	1,235.64
<i>Cedrus deodara</i>	274	0.89	5.56	948.64	25,687.43	229.54	0.10	1,097.80
<i>Acer rubrum</i>	258	0.84	1.08	184.18	3,106.44	27.76	0.01	132.76
<i>Cercis canadensis</i>	244	0.79	0.53	90.67	7,125.79	63.68	0.03	304.53
<i>Quercus agrifolia</i>	236	0.77	3.35	572.07	31,999.64	285.95	0.12	1,367.57
<i>Ulmus davidiana v. japonica</i>	232	0.76	1.05	179.84	3,207.57	28.66	0.01	137.08
<i>Olea europaea</i>	226	0.74	2.81	479.89	26,542.68	237.18	0.10	1,134.35
<i>Celtis occidentalis</i>	222	0.72	0.48	82.70	23,704.65	211.82	0.09	1,013.06
<i>Fraxinus uhdei</i>	214	0.70	4.45	759.39	29,569.67	264.23	0.11	1,263.72
<i>Zelkova serrata</i>	212	0.69	0.60	101.54	6,597.74	58.96	0.03	281.97
<i>Juglans hindsii</i>	201	0.65	5.19	885.71	27,836.74	248.75	0.11	1,189.66
<i>Laurus nobilis</i>	198	0.65	3.61	615.42	11,394.40	101.82	0.04	486.96
<i>Malus spp.</i>	192	0.63	1.11	189.81	2,432.22	21.73	0.01	103.95
<i>Quercus wislizeni</i>	189	0.62	2.88	490.52	22,964.37	205.21	0.09	981.43
<i>Prunus cerasifera</i>	188	0.61	1.82	310.81	8,957.85	80.05	0.03	382.83
<i>Arbutus unedo</i>	187	0.61	1.16	197.81	8,934.88	79.84	0.03	381.85
<i>Liriodendron tulipifera</i>	187	0.61	2.74	467.11	22,426.97	200.41	0.09	958.46

Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Avoided Runoff (gallon/yr.)	Avoided Runoff (\$/yr.)	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Quercus rubra</i>	183	0.60	0.36	61.18	2,459.74	21.98	0.01	105.12
<i>Eucalyptus sideroxylon</i>	178	0.58	1.46	248.32	40,520.63	362.09	0.15	1,731.73
<i>Acer x freemanii</i>	177	0.58	2.50	426.10	6,937.51	61.99	0.03	296.49
<i>Gymnocladus dioica</i>	176	0.57	1.71	292.45	6,399.06	57.18	0.02	273.48
<i>Morus alba</i>	172	0.56	2.27	386.57	22,328.83	199.53	0.08	954.27
<i>Fraxinus angustifolia ssp. Oxycarpa</i>	169	0.55	6.38	1,088.70	23,377.30	208.90	0.09	999.07
<i>Magnolia grandiflora</i>	160	0.52	1.39	236.76	9,031.91	80.71	0.03	386.00
<i>Pinus halepensis</i>	157	0.51	3.01	512.75	28,690.86	256.38	0.11	1,226.16
<i>Quercus buckleyi</i>	153	0.50	0.35	58.93	702.35	6.28	0.00	30.02
<i>Pistacia spp.</i>	152	0.50	0.11	19.36	400.66	3.58	0.00	17.12
<i>Quercus ilex</i>	152	0.50	4.21	717.90	32,154.57	287.33	0.12	1,374.19
<i>Ulmus spp.</i>	145	0.47	1.27	217.15	3,137.40	28.04	0.01	134.08
<i>Quercus canbyi</i>	143	0.47	0.24	41.56	355.40	3.18	0.00	15.19
<i>Melia azedarach</i>	125	0.41	3.71	633.27	16,478.61	147.25	0.06	704.25
<i>Fraxinus spp.</i>	124	0.40	2.41	411.20	10,310.63	92.14	0.04	440.64
<i>Prunus spp.</i>	122	0.40	0.96	163.92	5,377.52	48.05	0.02	229.82
<i>x Chitalpa tashkentensis</i>	113	0.37	0.48	81.51	1,171.08	10.46	0.00	50.05
<i>Tilia cordata</i>	95	0.31	0.98	166.73	7,461.37	66.67	0.03	318.88
<i>Sophora japonica</i>	90	0.29	0.15	26.04	12,134.75	108.44	0.05	518.60
<i>Quercus shumardii</i>	86	0.28	0.41	70.19	2,428.40	21.70	0.01	103.78
<i>Populus fremontii</i>	85	0.28	2.15	366.64	10,145.01	90.66	0.04	433.57
<i>Acer platanoides</i>	84	0.27	0.66	113.09	1,465.21	13.09	0.01	62.62
<i>Pinus brutia</i>	82	0.27	2.44	416.85	9,066.99	81.02	0.03	387.49
<i>Robinia pseudoacacia</i>	82	0.27	1.13	192.57	4,613.69	41.23	0.02	197.18
<i>Quercus coccinea</i>	80	0.26	0.83	142.01	4,212.51	37.64	0.02	180.03
<i>Ligustrum lucidum</i>	78	0.25	1.06	180.25	8,580.38	76.67	0.03	366.70
<i>Chilopsis linearis</i>	77	0.25	0.09	15.45	683.27	6.11	0.00	29.20
<i>Quercus robur</i>	77	0.25	0.46	78.76	2,231.66	19.94	0.01	95.37
<i>Prunus serrulata</i>	73	0.24	0.68	115.37	2,153.72	19.25	0.01	92.04
<i>Searsia lancea</i>	66	0.22	0.11	19.52	4,341.39	38.79	0.02	185.54
<i>Carpinus betulus</i>	64	0.21	0.54	91.68	6,984.04	62.41	0.03	298.48
<i>Juglans regia</i>	63	0.21	1.02	173.24	7,329.19	65.49	0.03	313.23
<i>Cupressus arizonica</i>	59	0.19	1.49	253.34	1,940.15	17.34	0.01	82.92
<i>Maclura pomifera</i>	56	0.18	0.03	4.79	104.28	0.93	0.00	4.46
<i>Pinus pinea</i>	55	0.18	1.42	242.81	11,508.48	102.84	0.04	491.84
<i>Ostrya virginiana</i>	52	0.17	0.27	45.48	2,756.95	24.64	0.01	117.82
<i>Populus nigra v. italica</i>	52	0.17	1.61	273.78	4,392.19	39.25	0.02	187.71
<i>Chionanthus retusus</i>	44	0.14	0.07	12.13	169.58	1.52	0.00	7.25
<i>Schinus molle</i>	44	0.14	0.16	26.48	3,241.61	28.97	0.01	138.54
<i>Acer campestre</i>	43	0.14	0.36	61.10	6,826.65	61.00	0.03	291.75
<i>Acacia melanoxylon</i>	41	0.13	0.84	143.26	7,321.71	65.43	0.03	312.91
<i>Salix spp.</i>	40	0.13	1.41	240.18	3,371.02	30.12	0.01	144.07
<i>Pinus nigra</i>	38	0.12	0.19	32.36	1,063.36	9.50	0.00	45.45

Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Avoided Runoff (gallon/yr.)	Avoided Runoff (\$/yr.)	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Eucalyptus camaldulensis</i>	36	0.12	0.27	45.75	15,278.44	136.53	0.06	652.95
<i>Liquidambar styraciflua</i>	33	0.11	0.46	78.64	4,268.68	38.15	0.02	182.43
<i>Acer saccharinum</i>	31	0.10	0.72	123.28	5,101.34	45.59	0.02	218.02
<i>Aesculus californica</i>	31	0.10	0.25	43.37	2,147.66	19.19	0.01	91.78
<i>Calocedrus decurrens</i>	31	0.10	0.24	41.27	2,952.49	26.38	0.01	126.18
<i>Albizia julibrissin</i>	30	0.10	0.14	24.54	2,999.84	26.81	0.01	128.20
<i>Chamaecyparis</i>	29	0.09	0.73	124.50	1,774.98	15.86	0.01	75.86
<i>Quercus macrocarpa</i>	29	0.09	0.46	78.16	5,251.85	46.93	0.02	224.45
<i>Geijera parviflora</i>	27	0.09	0.36	61.80	963.34	8.61	0.00	41.17
<i>Robinia spp.</i>	26	0.08	0.41	70.56	1,807.74	16.15	0.01	77.26
<i>Umbellularia californica</i>	26	0.08	0.75	128.38	3,051.30	27.27	0.01	130.40
<i>Chilopsis spp.</i>	25	0.08	0.03	4.33	41.89	0.37	0.00	1.79
<i>Platanus occidentalis</i>	24	0.08	0.41	70.44	5,027.43	44.93	0.02	214.86
<i>Fraxinus ornus</i>	23	0.07	0.16	27.44	3,581.37	32.00	0.01	153.06
<i>Prunus dulcis</i>	23	0.07	0.16	28.12	832.96	7.44	0.00	35.60
<i>Quercus spp.</i>	23	0.07	0.27	45.53	1,791.96	16.01	0.01	76.58
<i>Tilia tomentosa</i>	22	0.07	0.05	8.50	446.21	3.99	0.00	19.07
<i>Betula pendula</i>	21	0.07	0.22	37.76	972.17	8.69	0.00	41.55
<i>Quercus frainetto</i>	21	0.07	0.13	21.68	469.10	4.19	0.00	20.05
<i>Pinus thunbergii</i>	20	0.07	0.20	33.46	1,585.20	14.17	0.01	67.75
<i>Casuarina stricta</i>	19	0.06	1.45	248.00	1,685.51	15.06	0.01	72.03
<i>Parrotia persica</i>	19	0.06	0.01	1.33	48.92	0.44	0.00	2.09
<i>Tilia americana</i>	18	0.06	0.19	32.04	2,386.81	21.33	0.01	102.01
<i>Ulmus americana</i>	18	0.06	0.61	104.39	3,217.22	28.75	0.01	137.49
<i>Eucalyptus polyanthemos</i>	17	0.06	0.11	18.33	5,895.25	52.68	0.02	251.95
<i>Ficus spp.</i>	17	0.06	0.30	50.72	1,107.79	9.90	0.00	47.34
<i>Platanus orientalis</i>	17	0.06	0.21	35.42	2,158.50	19.29	0.01	92.25
<i>Cinnamomum camphora</i>	16	0.05	0.33	56.66	1,093.20	9.77	0.00	46.72
<i>Taxodium distichum</i>	16	0.05	0.01	1.69	19.97	0.18	0.00	0.85
<i>Catalpa speciosa</i>	15	0.05	0.13	22.19	1,125.82	10.06	0.00	48.11
<i>Prosopis chilensis</i>	15	0.05	0.01	2.26	14.51	0.13	0.00	0.62
<i>Prunus persica</i>	15	0.05	0.17	29.61	967.83	8.65	0.00	41.36
<i>Tilia x euchlora</i>	15	0.05	0.50	85.94	2,044.58	18.27	0.01	87.38
<i>Acer palmatum</i>	14	0.05	0.02	2.78	63.06	0.56	0.00	2.70
<i>Ceratonia siliqua</i>	14	0.05	0.02	3.69	1,948.45	17.41	0.01	83.27
<i>Juglans nigra</i>	14	0.05	0.26	44.47	2,520.64	22.52	0.01	107.72
<i>Maytenus boaria</i>	13	0.04	0.25	41.83	1,263.39	11.29	0.00	53.99
<i>Pinus sylvestris</i>	13	0.04	0.17	28.58	1,518.02	13.56	0.01	64.87
<i>Paulownia tomentosa</i>	12	0.04	0.07	12.31	594.40	5.31	0.00	25.40
<i>Tristanopsis spp.</i>	12	0.04	0.01	1.17	88.57	0.79	0.00	3.79
<i>Acer negundo</i>	11	0.04	0.08	13.68	402.00	3.59	0.00	17.18
<i>Afrocarpus gracilior</i>	11	0.04	0.07	11.98	771.32	6.89	0.00	32.96
<i>Alnus cordata</i>	11	0.04	0.12	20.99	1,432.07	12.80	0.01	61.20

Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Avoided Runoff (gallon/yr.)	Avoided Runoff (\$/yr.)	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Cedrus atlantica</i>	11	0.04	0.13	22.75	637.34	5.70	0.00	27.24
<i>Quercus cerris</i>	11	0.04	0.32	54.00	3,127.16	27.94	0.01	133.64
<i>Alnus rhombifolia</i>	10	0.03	0.06	10.92	539.64	4.82	0.00	23.06
<i>Eriobotrya japonica</i>	10	0.03	0.09	15.89	271.84	2.43	0.00	11.62
<i>Quercus palustris</i>	10	0.03	0.08	14.10	294.73	2.63	0.00	12.60
<i>Acer truncatum x platanoides</i>	9	0.03	0.18	29.95	669.73	5.98	0.00	28.62
<i>Eucalyptus melliodora</i>	9	0.03	0.11	18.65	3,910.67	34.95	0.01	167.13
<i>Prunus Kanzan</i>	9	0.03	0.08	12.80	263.09	2.35	0.00	11.24
<i>Pterocarya stenoptera</i>	9	0.03	0.13	21.94	1,322.71	11.82	0.01	56.53
<i>Punica spp.</i>	9	0.03	0.08	14.01	627.99	5.61	0.00	26.84
<i>Quercus laurifolia</i>	9	0.03	0.19	31.85	1,112.95	9.95	0.00	47.56
<i>Ulmus procera</i>	9	0.03	0.33	56.61	1,353.75	12.10	0.01	57.86
<i>Magnolia x soulangeana</i>	8	0.03	0.04	6.25	252.69	2.26	0.00	10.80
<i>Melaleuca linariifolia</i>	8	0.03	0.01	0.86	741.99	6.63	0.00	31.71
<i>Ailanthus altissima</i>	7	0.02	0.14	24.46	532.09	4.75	0.00	22.74
<i>Crataegus spp.</i>	7	0.02	0.04	7.19	175.04	1.56	0.00	7.48
<i>Eucalyptus spp.</i>	7	0.02	0.03	4.32	4,030.43	36.02	0.02	172.25
<i>Quercus chrysolepis</i>	7	0.02	-	0.25	9.13	0.08	0.00	0.39
<i>Acer pseudoplatanus</i>	6	0.02	0.10	16.83	836.47	7.47	0.00	35.75
<i>Eucalyptus nicholii</i>	6	0.02	0.05	8.06	2,188.13	19.55	0.01	93.51
<i>Quercus muehlenbergii</i>	6	0.02	0.04	7.09	166.59	1.49	0.00	7.12
<i>Salix babylonica</i>	6	0.02	0.27	46.12	1,270.19	11.35	0.00	54.28
<i>Betula nigra</i>	5	0.02	0.02	2.56	61.64	0.55	0.00	2.63
<i>Carpinus caroliniana</i>	5	0.02	0.02	2.92	141.98	1.27	0.00	6.07
<i>Quercus imbricaria</i>	5	0.02	0.01	1.45	8.68	0.08	0.00	0.37
<i>Xylosma spp.</i>	5	0.02	0.05	9.23	146.24	1.31	0.00	6.25
<i>Cupressus sempervirens</i>	4	0.01	0.03	5.38	205.49	1.84	0.00	8.78
<i>Lagerstroemia subcostata</i>	4	0.01	-	0.73	7.63	0.07	0.00	0.33
<i>Magnoliopsida</i>	4	0.01	-	0.57	5.54	0.05	0.00	0.24
<i>Myrtuss pp.</i>	4	0.01	0.01	1.25	1,256.13	11.22	0.00	53.68
<i>Picea pungens</i>	4	0.01	0.02	2.75	115.80	1.03	0.00	4.95
<i>Thuja plicata</i>	4	0.01	0.01	1.84	386.22	3.45	0.00	16.51
<i>Ulmus pumila</i>	4	0.01	0.20	33.75	660.16	5.90	0.00	28.21
<i>Acer griseum</i>	3	0.01	0.01	2.37	52.14	0.47	0.00	2.23
<i>Brachychiton populneus</i>	3	0.01	0.05	7.86	283.59	2.53	0.00	12.12
<i>Eucalyptus globulus</i>	3	0.01	0.03	4.52	1,292.93	11.55	0.00	55.26
<i>Juglans spp.</i>	3	0.01	0.06	10.51	264.81	2.37	0.00	11.32
<i>Metasequoia glyptostroboides</i>	3	0.01	-	0.23	3.29	0.03	0.00	0.14
<i>Parkinsonia aculeata</i>	3	0.01	0.01	0.86	12.34	0.11	0.00	0.53
<i>Pinus radiata</i>	3	0.01	0.05	9.35	397.29	3.55	0.00	16.98
<i>Acer spp.</i>	2	0.01	0.06	9.40	202.12	1.81	0.00	8.64
<i>Acer saccharum</i>	2	0.01	0.04	6.13	353.45	3.16	0.00	15.10

Species	Number of Trees	% of Pop.	Carbon Sequestration (ton/yr.)	Carbon Sequestration (\$/yr.)	Avoided Runoff (gallon/yr.)	Avoided Runoff (\$/yr.)	Pollution Removal (ton/yr.)	Pollution Removal (\$/yr.)
<i>Callistemon spp.</i>	2	0.01	-	0.11	6.21	0.06	0.00	0.27
<i>Citrus spp.</i>	2	0.01	0.01	1.00	129.04	1.15	0.00	5.52
<i>Citrus limon</i>	2	0.01	0.02	3.57	19.67	0.18	0.00	0.84
<i>Corymbia maculata</i>	2	0.01	0.01	2.14	503.36	4.50	0.00	21.51
<i>Crataegus phaenopyrum</i>	2	0.01	-	0.66	5.76	0.05	0.00	0.25
<i>Koelreuteria elegans</i>	2	0.01	0.01	2.34	69.27	0.62	0.00	2.96
<i>Photinia spp.</i>	2	0.01	0.01	2.55	26.56	0.24	0.00	1.14
<i>Pinus coulteri</i>	2	0.01	0.02	3.76	84.75	0.76	0.00	3.62
<i>Prunus armeniaca</i>	2	0.01	0.02	3.54	39.35	0.35	0.00	1.68
<i>Prunus lyonii</i>	2	0.01	0.03	4.46	58.27	0.52	0.00	2.49
<i>Prunus x yedoensis</i>	2	0.01	0.01	0.86	11.22	0.10	0.00	0.48
<i>Quercus castaneifolia</i>	2	0.01	0.01	2.11	28.28	0.25	0.00	1.21
<i>Quercus nigra</i>	2	0.01	0.02	3.66	113.18	1.01	0.00	4.84
<i>Quercus phellos</i>	2	0.01	0.06	10.29	579.07	5.17	0.00	24.75
<i>Washingtonia spp.</i>	2	0.01	-	0.81	58.87	0.53	0.00	2.52
<i>Acer macrophyllum</i>	1	0.00	-	0.26	2.84	0.03	0.00	0.12
<i>Brachychiton acerifolius</i>	1	0.00	-	0.08	2.17	0.02	0.00	0.09
<i>Carya illinoensis</i>	1	0.00	0.01	1.36	88.87	0.79	0.00	3.80
<i>Cercidiphyllum japonicum</i>	1	0.00	-	0.05	3.44	0.03	0.00	0.15
<i>Elaeocarpus decipiens</i>	1	0.00	-	0.21	2.99	0.03	0.00	0.13
<i>Firmiana simplex</i>	1	0.00	0.01	1.79	50.42	0.45	0.00	2.15
<i>Juniperus virginiana</i>	1	0.00	0.01	1.06	55.13	0.49	0.00	2.36
<i>Larix occidentalis</i>	1	0.00	0.01	0.93	19.30	0.17	0.00	0.83
<i>Magnolia virginiana</i>	1	0.00	0.01	2.53	105.55	0.94	0.00	4.51
<i>Malus x arnoldiana</i>	1	0.00	-	-	-	-	0.00	-
<i>Nyssa sylvatica</i>	1	0.00	-	0.07	0.75	0.01	0.00	0.03
<i>Ostrya carpinifolia</i>	1	0.00	-	0.30	5.61	0.05	0.00	0.24
<i>Persea spp.</i>	1	0.00	-	0.33	4.26	0.04	0.00	0.18
<i>Pistacia vera</i>	1	0.00	0.01	1.54	35.31	0.32	0.00	1.51
<i>Prunus subhirtella</i>	1	0.00	-	0.27	152.23	1.36	0.00	6.51
<i>Prunus x blireiana</i>	1	0.00	0.02	2.67	72.49	0.65	0.00	3.10
<i>Pyrus betulifolia</i>	1	0.00	0.01	1.29	16.16	0.14	0.00	0.69
<i>Quercus engelmannii</i>	1	0.00	0.02	3.61	188.96	1.69	0.00	8.08
<i>Tetradium spp.</i>	1	0.00	-	0.19	48.85	0.44	0.00	2.09
<i>Thuja spp.</i>	1	0.00	0.06	10.39	267.50	2.39	0.00	11.43
<i>Ulmus x hollandica</i>	1	0.00	-	0.57	263.91	2.36	0.00	11.28
<i>Washingtonia filifera</i>	1	0.00	-	0.12	0.97	0.01	0.00	0.04